



Action myoclonus–renal failure syndrome

Action myoclonus–renal failure (AMRF) syndrome causes episodes of involuntary muscle jerking or twitching (myoclonus) and, often, kidney (renal) disease. Although the condition name refers to kidney disease, not everyone with the condition has problems with kidney function.

The movement problems associated with AMRF syndrome typically begin with involuntary rhythmic shaking (tremor) in the fingers and hands that occurs at rest and is most noticeable when trying to make small movements, such as writing. Over time, tremors can affect other parts of the body, such as the head, torso, legs, and tongue. Eventually, the tremors worsen to become myoclonic jerks, which can be triggered by voluntary movements or the intention to move (action myoclonus). These myoclonic jerks typically occur in the torso; upper and lower limbs; and face, particularly the muscles around the mouth and the eyelids. Anxiety, excitement, stress, or extreme tiredness (fatigue) can worsen the myoclonus. Some affected individuals develop seizures, a loss of sensation and weakness in the limbs (peripheral neuropathy), or hearing loss caused by abnormalities in the inner ear (sensorineural hearing loss). Severe seizures or myoclonus can be life-threatening.

When kidney problems occur, an early sign is excess protein in the urine (proteinuria). Kidney function worsens over time, until the kidneys are no longer able to filter fluids and waste products from the body effectively (end-stage renal disease).

AMRF syndrome typically begins causing symptoms between ages 15 and 25, but it can appear at younger or older ages. The age of onset and the course of the condition vary, even among members of the same family. Either the movement problems or kidney disease can occur first, or they can begin at the same time. Most people survive 7 to 15 years after the symptoms appear.

Frequency

AMRF syndrome is a rare condition that has been found worldwide. Its exact prevalence is unknown. At least 38 individuals with the condition have been described in the medical literature.

Causes

AMRF syndrome is caused by mutations in the *SCARB2* gene. This gene provides instructions for making the LIMP-2 protein, which transports an enzyme called beta-glucocerebrosidase to cellular structures called lysosomes. Lysosomes are specialized compartments that digest and recycle materials. In these compartments, beta-glucocerebrosidase breaks down a fatty substance called glucocerebroside. The

LIMP-2 protein remains in the lysosome after transporting beta-glucocerebrosidase and is important for the stability of these structures.

SCARB2 gene mutations associated with AMRF syndrome lead to production of an altered LIMP-2 protein that cannot get to the lysosome. As a result, the movement of beta-glucocerebrosidase to lysosomes is impaired. It is thought that a shortage of beta-glucocerebrosidase function in these structures contributes to the signs and symptoms of AMRF syndrome, although the mechanism is unclear. Researchers are working to understand why some people with *SCARB2* gene mutations have kidney problems and others do not.

Inheritance Pattern

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

Other Names for This Condition

- action myoclonus-renal failure syndrome
- action myoclonus–renal failure syndrome
- AMRF
- epilepsy, progressive myoclonic 4, with or without renal failure
- EPM4
- familial myoclonus with renal failure
- myoclonus-nephropathy syndrome
- progressive myoclonus epilepsy with renal failure

Diagnosis & Management

Genetic Testing Information

- What is genetic testing?
[/primer/testing/genetic-testing](#)
- Genetic Testing Registry: Epilepsy, progressive myoclonic 4, with or without renal failure
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0751779/>

Other Diagnosis and Management Resources

- GeneReview: Action Myoclonus - Renal Failure Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK333437>

Additional Information & Resources

Health Information from MedlinePlus

- Health Topic: Kidney Diseases
<https://medlineplus.gov/kidneydiseases.html>
- Health Topic: Movement Disorders
<https://medlineplus.gov/movementdisorders.html>

Additional NIH Resources

- National Institute of Neurological Disorders and Stroke: Myoclonus Fact Sheet
<https://www.ninds.nih.gov/Disorders/All-Disorders/Myoclonus-Information-Page>

Educational Resources

- Baylor College of Medicine: Myoclonus
<https://www.bcm.edu/healthcare/care-centers/parkinsons/conditions/myoclonus>
- MalaCards: epilepsy, progressive myoclonic, 4, with or without renal failure
https://www.malacards.org/card/epilepsy_progressive_myoclonic_4_with_or_without_renal_failure_2
- Merck Manual Consumer Version: Myoclonus
<https://www.merckmanuals.com/home/brain-spinal-cord-and-nerve-disorders/movement-disorders/myoclonus>
- Orphanet: Action myoclonus-renal failure syndrome
https://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=163696

Patient Support and Advocacy Resources

- American Epilepsy Society
<https://www.aesnet.org/>
- American Kidney Fund
<http://www.kidneyfund.org/>
- Citizens United for Research in Epilepsy (CURE)
<https://www.cureepilepsy.org/>
- National Kidney Foundation
<https://www.kidney.org/>

Clinical Information from GeneReviews

- Action Myoclonus - Renal Failure Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK333437>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28Myoclonic+Epilepsies,+Progressive%5BMAJR%5D%29+AND+%28action+myoclonus-renal+failure+syndrome%5BTIAB%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

Catalog of Genes and Diseases from OMIM

- EPILEPSY, PROGRESSIVE MYOCLONIC, 4, WITH OR WITHOUT RENAL FAILURE
<http://omim.org/entry/254900>

Sources for This Summary

- Amrom D, Andermann F, Andermann E. Action Myoclonus – Renal Failure Syndrome. 2015 Dec 17. In: Pagon RA, Adam MP, Ardinger HH, Wallace SE, Amemiya A, Bean LJH, Bird TD, Ledbetter N, Mefford HC, Smith RJH, Stephens K, editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2017. Available from <http://www.ncbi.nlm.nih.gov/books/NBK333437/>
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/26677510>
- Balreira A, Gaspar P, Caiola D, Chaves J, Beirão I, Lima JL, Azevedo JE, Miranda MC. A nonsense mutation in the LIMP-2 gene associated with progressive myoclonic epilepsy and nephrotic syndrome. *Hum Mol Genet.* 2008 Jul 15;17(14):2238-43. doi: 10.1093/hmg/ddn124. Epub 2008 Apr 17.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18424452>
- Berkovic SF, Dibbens LM, Oshlack A, Silver JD, Katerelos M, Vears DF, Lüllmann-Rauch R, Blanz J, Zhang KW, Stankovich J, Kalnins RM, Dowling JP, Andermann E, Andermann F, Faldini E, D'Hooge R, Vadlamudi L, Macdonell RA, Hodgson BL, Bayly MA, Savige J, Mulley JC, Smyth GK, Power DA, Saftig P, Bahlo M. Array-based gene discovery with three unrelated subjects shows SCARB2/LIMP-2 deficiency causes myoclonus epilepsy and glomerulosclerosis. *Am J Hum Genet.* 2008 Mar;82(3):673-84. doi: 10.1016/j.ajhg.2007.12.019. Epub 2008 Feb 28.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18308289>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2427287/>
- Blanz J, Groth J, Zachos C, Wehling C, Saftig P, Schwake M. Disease-causing mutations within the lysosomal integral membrane protein type 2 (LIMP-2) reveal the nature of binding to its ligand beta-glucocerebrosidase. *Hum Mol Genet.* 2010 Feb 15;19(4):563-72. doi: 10.1093/hmg/ddp523. Epub 2009 Nov 20.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19933215>

- Dibbens LM, Michelucci R, Gambardella A, Andermann F, Rubboli G, Bayly MA, Joensuu T, Vears DF, Franceschetti S, Canafoglia L, Wallace R, Bassuk AG, Power DA, Tassinari CA, Andermann E, Lehesjoki AE, Berkovic SF. SCARB2 mutations in progressive myoclonus epilepsy (PME) without renal failure. *Ann Neurol.* 2009 Oct;66(4):532-6. doi: 10.1002/ana.21765.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19847901>
 - Hopfner F, Schormair B, Knauf F, Berthele A, Tölle TR, Baron R, Maier C, Treede RD, Binder A, Sommer C, Maihöfner C, Kunz W, Zimprich F, Heemann U, Pfeufer A, Näbauer M, Kääh S, Nowak B, Gieger C, Lichtner P, Trenkwalder C, Oexle K, Winkelmann J. Novel SCARB2 mutation in action myoclonus-renal failure syndrome and evaluation of SCARB2 mutations in isolated AMRF features. *BMC Neurol.* 2011 Oct 27;11:134. doi: 10.1186/1471-2377-11-134.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22032306>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3222607/>
-

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/action-myoclonus-renal-failure-syndrome>

Reviewed: June 2016

Published: June 11, 2019

Lister Hill National Center for Biomedical Communications

U.S. National Library of Medicine

National Institutes of Health

Department of Health & Human Services