



Branchio-oculo-facial syndrome

Branchio-oculo-facial syndrome is a condition that affects development before birth, particularly of structures in the face and neck. Its characteristic features include skin anomalies on the neck, malformations of the eyes and ears, and distinctive facial features.

"Branchio-" refers to the branchial arches, which are structures in the developing embryo that give rise to tissues in the face and neck. In people with branchio-oculo-facial syndrome, the first and second branchial arches do not develop properly, leading to abnormal patches of skin, typically on the neck or near the ears. These patches can be unusually thin, hairy, or red and densely packed with blood vessels (hemangiomas). In a small number of individuals, tissue from a gland called the thymus is abnormally located on the skin of the neck (dermal thymus). Problems with branchial arch development underlie many of the other features of branchio-oculo-facial syndrome.

"Oculo-" refers to the eyes. Many people with branchio-oculo-facial syndrome have malformations of the eyes that can lead to vision impairment. These abnormalities include unusually small eyeballs (microphthalmia), no eyeballs (anophthalmia), a gap or split in structures that make up the eyes (coloboma), or blockage of the tear ducts (nasolacrimal duct stenosis).

Problems with development of the face lead to distinctive facial features in people with branchio-oculo-facial syndrome. Many affected individuals have a split in the upper lip (cleft lip) or a pointed upper lip that resembles a poorly repaired cleft lip (often called a pseudocleft lip) with or without an opening in the roof of the mouth (cleft palate). Other facial characteristics include widely spaced eyes (hypertelorism), an increased distance between the inner corners of the eyes (telecanthus), outside corners of the eyes that point upward (upslanting palpebral fissures), a broad nose with a flattened tip, and weakness of the muscles in the lower face. The ears are also commonly affected, resulting in malformed or prominent ears. Abnormalities of the inner ear or of the tiny bones in the ears (ossicles) can cause hearing loss in people with this condition.

Branchio-oculo-facial syndrome can affect other structures and tissues as well. Some affected individuals have kidney abnormalities, such as malformed kidneys or multiple kidney cysts. Nail and teeth abnormalities also occur, and some people with this condition have prematurely graying hair.

Frequency

Branchio-oculo-facial syndrome is a rare condition, although the prevalence is unknown.

Causes

Branchio-oculo-facial syndrome is caused by mutations in the *TFAP2A* gene. This gene provides instructions for making a protein called transcription factor AP-2 alpha (AP-2 α). As its name suggests, this protein is a transcription factor, which means it attaches (binds) to specific regions of DNA and helps control the activity of particular genes. Transcription factor AP-2 α regulates genes that are involved in several cellular processes, such as cell division and the self-destruction of cells that are no longer needed (apoptosis). This protein is critical during development before birth, particularly of the branchial arches, which form the structures of the face and neck.

Most *TFAP2A* gene mutations that cause branchio-oculo-facial syndrome change single protein building blocks (amino acids) in the transcription factor AP-2 α protein. These changes tend to occur in a region of the protein that allows it to bind to DNA. Without this function, transcription factor AP-2 α cannot control the activity of genes during development, which disrupts the development of the eyes, ears, and face and causes the features of branchio-oculo-facial syndrome.

Inheritance Pattern

Branchio-oculo-facial syndrome is inherited in an autosomal dominant pattern, which means one copy of the altered gene in each cell is sufficient to cause the disorder.

In about half of cases, an affected person inherits the mutation from one affected parent. The remaining cases occur in people whose parents do not have a mutation in the *TFAP2A* gene. In these situations, the mutation likely occurs as a random event during the formation of reproductive cells (eggs and sperm) in a parent or in early fetal development of the affected individual.

Other Names for This Condition

- BOFS
- branchial clefts with characteristic facies, growth retardation, imperforate nasolacrimal duct, and premature aging
- hemangiomatous branchial clefts-lip pseudocleft syndrome
- lip pseudocleft-hemangiomatous branchial cyst syndrome

Diagnosis & Management

Genetic Testing Information

- What is genetic testing?
[/primer/testing/genetictesting](#)
- Genetic Testing Registry: Branchiooculofacial syndrome
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0376524/>

Research Studies from ClinicalTrials.gov

- ClinicalTrials.gov
<https://clinicaltrials.gov/ct2/results?cond=%22branchio-oculo-facial+syndrome%22+OR+%22Craniofacial+Abnormalities%22>

Other Diagnosis and Management Resources

- GeneReview: Branchiooculofacial Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK55063>
- MedlinePlus Encyclopedia: Cleft Lip and Palate
<https://medlineplus.gov/ency/article/001051.htm>

Additional Information & Resources

Health Information from MedlinePlus

- Encyclopedia: Cleft Lip and Palate
<https://medlineplus.gov/ency/article/001051.htm>
- Health Topic: Cleft Lip and Palate
<https://medlineplus.gov/cleftlipandpalate.html>
- Health Topic: Craniofacial Abnormalities
<https://medlineplus.gov/craniofacialabnormalities.html>

Genetic and Rare Diseases Information Center

- Branchiooculofacial syndrome
<https://rarediseases.info.nih.gov/diseases/3212/branchiooculofacial-syndrome>

Educational Resources

- American Association for Pediatric Ophthalmology and Strabismus: Coloboma
<https://aapos.org/browse/glossary/entry?GlossaryKey=ae5e1cdf-4e2e-4e35-8350-1ac447506955>
- KidsHealth from Nemours: Cleft Lip and Palate
<https://kidshealth.org/en/parents/cleft-lip-palate.html>
- March of Dimes: Cleft Lip and Cleft Palate
<https://www.marchofdimes.org/baby/cleft-lip-and-cleft-palate.aspx>
- Orphanet: Branchio-oculo-facial syndrome
https://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=1297
- The University of Arizona Health Sciences
<https://disorders.eyes.arizona.edu/disorders/branchiooculofacial-syndrome>

Patient Support and Advocacy Resources

- About Face
<https://www.aboutface.ca/>
- Ameriface
<http://www.ameriface.org/>
- National Organization for Rare Disorders (NORD)
<https://rarediseases.org/rare-diseases/branchio-oculo-facial-syndrome/>

Clinical Information from GeneReviews

- Branchiooculofacial Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK55063>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28branchio-oculo-facial+syndrome%5BTIAB%5D%29+OR+%28branchiooculofacial+syndrome%5BTIAB%5D%29%29+AND+english%5BIa%5D+AND+human%5Bmh%5D+AND+%22last+2880+days%22%5Bdp%5D>

Catalog of Genes and Diseases from OMIM

- BRANCHIOOCULOFACIAL SYNDROME
<http://omim.org/entry/113620>

Sources for This Summary

- Developmental Biology (sixth edition, 2000): The Neural Crest
<https://www.ncbi.nlm.nih.gov/books/NBK10065/>
- Lin AE, Gorlin RJ, Lurie IW, Brunner HG, van der Burgt I, Naumchik IV, Rummyantseva NV, Stengel-Rutkowski S, Rosenbaum K, Meinecke P, et al. Further delineation of the branchio-oculo-facial syndrome. *Am J Med Genet.* 1995 Mar 13;56(1):42-59. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/7747785>
- Lin AE, Milunsky JM. Branchiooculofacial Syndrome. 2011 May 31. In: Pagon RA, Adam MP, Ardinger HH, Wallace SE, Amemiya A, Bean LJH, Bird TD, Ledbetter N, Mefford HC, Smith RJH, Stephens K, editors. *GeneReviews®* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2017. Available from <http://www.ncbi.nlm.nih.gov/books/NBK55063/>
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21634087>

- Milunsky JM, Maher TA, Zhao G, Roberts AE, Stalker HJ, Zori RT, Burch MN, Clemens M, Mulliken JB, Smith R, Lin AE. TFAP2A mutations result in branchio-oculo-facial syndrome. *Am J Hum Genet.* 2008 May;82(5):1171-7. doi: 10.1016/j.ajhg.2008.03.005. Erratum in: *Am J Hum Genet.* 2009 Feb; 84(2):301..
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18423521>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2427243/>
 - Milunsky JM, Maher TM, Zhao G, Wang Z, Mulliken JB, Chitayat D, Clemens M, Stalker HJ, Bauer M, Burch M, Chénier S, Cunningham ML, Drack AV, Janssens S, Karlea A, Klatt R, Kini U, Klein O, Lachmeijer AM, Megarbane A, Mendelsohn NJ, Meschino WS, Mortier GR, Parkash S, Ray CR, Roberts A, Roberts A, Reardon W, Schnur RE, Smith R, Splitt M, Tezcan K, Whiteford ML, Wong DA, Zori R, Lin AE. Genotype-phenotype analysis of the branchio-oculo-facial syndrome. *Am J Med Genet A.* 2011 Jan;155A(1):22-32. doi: 10.1002/ajmg.a.33783.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21204207>
-

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/branchio-oculo-facial-syndrome>

Reviewed: September 2012

Published: May 14, 2019

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services