



Congenital anomalies of kidney and urinary tract

Congenital anomalies of kidney and urinary tract (CAKUT) is a group of abnormalities affecting the kidneys or other structures of the urinary tract. The additional parts of the urinary tract that may be affected include the bladder, the tubes that carry urine from each kidney to the bladder (the ureters), and the tube that carries urine from the bladder out of the body (the urethra). CAKUT results from abnormal development of the urinary system and is present from birth (congenital), although the abnormality may not become apparent until later in life.

Individuals with CAKUT have one or more kidney or urinary tract abnormalities. For paired structures, like the kidneys and ureters, one or both may be affected. Many different developmental abnormalities are classified as CAKUT, including underdevelopment or absence of a kidney (renal hypodysplasia or agenesis), a kidney formed of fluid-filled sacs called cysts (multicystic dysplastic kidney), buildup of urine in the kidneys (hydronephrosis), an extra ureter leading to the kidney (duplex kidney or duplicated collecting system), a blockage in a ureter where it joins the kidney (ureteropelvic junction obstruction), an abnormally wide ureter (megaureter), backflow of urine from the bladder into the ureter (vesicoureteral reflux), and an abnormal membrane in the urethra that blocks the flow of urine out of the bladder (posterior urethral valve).

CAKUT varies in severity. The abnormalities can result in recurrent urinary tract infections or a buildup of urine in the urinary tract, which may damage the kidneys or other structures. Severe CAKUT can result in life-threatening kidney failure and end-stage renal disease.

CAKUT is often one of several features of a condition that affects multiple body systems (syndromic CAKUT). For example, renal coloboma syndrome, 17q12 deletion syndrome, renal cysts and diabetes (RCAD) syndrome, Fraser syndrome, Townes-Brocks syndrome, and branchio-oto-renal syndrome can cause kidney or urinary tract abnormalities in addition to other problems. However, urinary system abnormalities sometimes occur without other signs and symptoms, which is known as nonsyndromic or isolated CAKUT.

Frequency

CAKUT is estimated to occur in 1 in 100 to 500 newborns. These abnormalities are the most common cause of end-stage renal disease in children.

Causes

The causes of CAKUT are complex. It is likely that a combination of genetic and environmental factors contribute to the formation of kidney and urinary tract abnormalities.

The genetic factors involved in most cases of CAKUT are unknown. Syndromic CAKUT is caused by changes in the genes associated with the particular syndrome. Variations in these same genes can also underlie some cases of isolated CAKUT. The genes most commonly associated with isolated CAKUT are *PAX2*, which is also associated with renal coloboma syndrome, and *HNF1B*, which is involved in 17q12 deletion syndrome and RCAD syndrome. These two genes play critical roles in the formation of the kidneys, urinary tract, and other tissues and organs during embryonic development. Certain mutations in these genes are thought to disrupt development of the kidneys or other parts of the urinary tract before birth, leading to CAKUT. Mutations in many other genes involved in development of the urinary system have also been associated with isolated or syndromic CAKUT.

Research shows that the same genetic mutation can lead to different kidney or urinary tract abnormalities, even among members of the same family. It is likely that additional changes in other genes help determine how the condition develops and how severe it is. In addition, environmental factors may influence development of CAKUT. The risk of CAKUT is higher in babies whose mothers had diabetes; took certain medications that are harmful to the kidneys, such as some anti-seizure drugs; or lacked certain vitamins and minerals, such as folate and iron, during pregnancy.

Inheritance Pattern

Inheritance of CAKUT is complex and not completely understood. About 10 to 20 percent of cases are thought to occur in families. When inherited, CAKUT most commonly follows an autosomal dominant pattern, which means one copy of the altered gene in each cell is sufficient to cause an abnormality. However, some people who have the altered gene never develop CAKUT, a situation known as reduced penetrance.

Less commonly, CAKUT follows an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not have signs or symptoms of the condition.

In many cases, the inheritance pattern is unknown or the condition is not inherited. In some of these cases, a new (de novo) mutation in the gene that occurs during the formation of reproductive cells (eggs or sperm) in an affected individual's parent or in early embryonic development may underlie the urinary system abnormality. These cases occur in people with no history of the disorder in their family.

Other Names for This Condition

- CAKUT

Diagnosis & Management

Genetic Testing Information

- What is genetic testing?
[/primer/testing/geneticTesting](#)
- Genetic Testing Registry: Congenital anomalies of kidney and urinary tract
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C196894/>

Research Studies from ClinicalTrials.gov

- ClinicalTrials.gov
<https://clinicaltrials.gov/ct2/results?cond=%22congenital+anomalies+of+kidney+and+urinary+tract%22+OR+%22CAKUT%22>

Other Diagnosis and Management Resources

- MedlinePlus Health Topic: Kidney Tests
<https://medlineplus.gov/kidneytests.html>
- National Institute of Diabetes and Digestive and Kidney Diseases: Urinary Tract Imaging
<https://www.niddk.nih.gov/health-information/diagnostic-tests/urinary-tract-imaging>
- National Kidney Foundation: Detecting Kidney and Urinary Tract Abnormalities Before Birth
<https://www.kidney.org/atoz/content/detectkid>

Additional Information & Resources

Health Information from MedlinePlus

- Encyclopedia: Urinary Tract Infection - Children
<https://medlineplus.gov/ency/article/000505.htm>
- Health Topic: Kidney Diseases
<https://medlineplus.gov/kidneydiseases.html>
- Health Topic: Kidney Tests
<https://medlineplus.gov/kidneytests.html>
- Health Topic: Ureteral Disorders
<https://medlineplus.gov/ureteraldisorders.html>
- Health Topic: Urethral Disorders
<https://medlineplus.gov/urethraldisorders.html>

Additional NIH Resources

- National Institute of Diabetes and Digestive and Kidney Diseases: The Urinary Tract and How It Works
<https://www.niddk.nih.gov/health-information/urologic-diseases/urinary-tract-how-it-works>

Educational Resources

- Boston Children's Hospital: Hydronephrosis
<http://www.childrenshospital.org/conditions-and-treatments/conditions/h/hydronephrosis>
- Children's Hospital of Philadelphia: Duplex Kidney
<https://www.chop.edu/conditions-diseases/duplex-kidney>
- Children's Hospital of Philadelphia: Multicystic Dysplastic Kidney
<https://www.chop.edu/conditions-diseases/multicystic-dysplastic-kidney>
- Children's Hospital of Philadelphia: Posterior Urethral Valves
<https://www.chop.edu/conditions-diseases/posterior-urethral-valves-puv>
- Children's Hospital of Philadelphia: Ureteropelvic Junction Obstruction
<https://www.chop.edu/conditions-diseases/ureteropelvic-junction-upj-obstruction>
- Children's National Health System: Megaureter
<https://childrensnational.org/choose-childrens/conditions-and-treatments/urology/megaureter>
- KidsHealth from Nemours: Kidney and Urinary Tract
<https://kidshealth.org/en/parents/kidneys-urinary.html>
- MalaCards: congenital anomalies of kidney and urinary tract 1
https://www.malacards.org/card/congenital_anomalies_of_kidney_and_urinary_tract_1
- MalaCards: congenital anomalies of kidney and urinary tract 2
https://www.malacards.org/card/congenital_anomalies_of_kidney_and_urinary_tract_2
- National Institute of Diabetes and Digestive and Kidney Diseases: Vesicoureteral Reflux
<https://www.niddk.nih.gov/health-information/urologic-diseases/urine-blockage-newborns/vesicoureteral-reflux>
- Nationwide Children's Hospital: Multicystic Dysplastic Kidney
<https://www.nationwidechildrens.org/conditions/multicystic-dysplastic-kidney>

Patient Support and Advocacy Resources

- March of Dimes
<https://www.marchofdimes.org/baby/genital-and-urinary-tract-defects.aspx>
- National Kidney Foundation
<https://www.kidney.org/>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28congenital+anomalies+of+kidney+and+urinary+tract%5BTIAB%5D%29+OR+%28CAKUT%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1440+days%22%5Bdp%5D>

Catalog of Genes and Diseases from OMIM

- CONGENITAL ANOMALIES OF KIDNEY AND URINARY TRACT 1
<http://omim.org/entry/610805>
- CONGENITAL ANOMALIES OF KIDNEY AND URINARY TRACT 2
<http://omim.org/entry/143400>

Medical Genetics Database from MedGen

- Congenital anomalies of kidney and urinary tract
<https://www.ncbi.nlm.nih.gov/medgen/369894>

Sources for This Summary

- Capone VP, Morello W, Taroni F, Montini G. Genetics of Congenital Anomalies of the Kidney and Urinary Tract: The Current State of Play. *Int J Mol Sci.* 2017 Apr 11;18(4). pii: E796. doi: 10.3390/ijms18040796. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/28398236>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5412380/>
- Hwang DY, Dworschak GC, Kohl S, Saisawat P, Vivante A, Hilger AC, Reutter HM, Soliman NA, Bogdanovic R, Kehinde EO, Tasic V, Hildebrandt F. Mutations in 12 known dominant disease-causing genes clarify many congenital anomalies of the kidney and urinary tract. *Kidney Int.* 2014 Jun;85(6):1429-33. doi: 10.1038/ki.2013.508. Epub 2014 Jan 15.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/24429398>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4040148/>
- Nicolaou N, Renkema KY, Bongers EM, Giles RH, Knoers NV. Genetic, environmental, and epigenetic factors involved in CAKUT. *Nat Rev Nephrol.* 2015 Dec;11(12):720-31. doi: 10.1038/rrneph.2015.140. Epub 2015 Aug 18. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/26281895>
- Sanna-Cherchi S, Westland R, Ghiggeri GM, Gharavi AG. Genetic basis of human congenital anomalies of the kidney and urinary tract. *J Clin Invest.* 2018 Jan 2;128(1):4-15. doi: 10.1172/JCI95300. Epub 2018 Jan 2. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/29293093>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5749511/>

- Vivante A, Kohl S, Hwang DY, Dworschak GC, Hildebrandt F. Single-gene causes of congenital anomalies of the kidney and urinary tract (CAKUT) in humans. *Pediatr Nephrol.* 2014 Apr;29(4): 695-704. doi: 10.1007/s00467-013-2684-4. Epub 2014 Jan 8. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/24398540>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4676405/>
 - Weber S, Moriniere V, Knüppel T, Charbit M, Dusek J, Ghiggeri GM, Jankauskienė A, Mir S, Montini G, Peco-Antic A, Wühl E, Zurowska AM, Mehls O, Antignac C, Schaefer F, Salomon R. Prevalence of mutations in renal developmental genes in children with renal hypodysplasia: results of the ESCAPE study. *J Am Soc Nephrol.* 2006 Oct;17(10):2864-70. Epub 2006 Sep 13.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16971658>
-

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/congenital-anomalies-of-kidney-and-urinary-tract>

Reviewed: November 2018

Published: December 11, 2018

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services