



## Familial exudative vitreoretinopathy

Familial exudative vitreoretinopathy is a hereditary disorder that can cause progressive vision loss. This condition affects the retina, the specialized light-sensitive tissue that lines the back of the eye. The disorder prevents blood vessels from forming at the edges of the retina, which reduces the blood supply to this tissue.

The signs and symptoms of familial exudative vitreoretinopathy vary widely, even within the same family. In many affected individuals, the retinal abnormalities never cause any vision problems. In others, a reduction in the retina's blood supply causes the retina to fold, tear, or separate from the back of the eye (retinal detachment). This retinal damage can lead to vision loss and blindness. Other eye abnormalities are also possible, including eyes that do not look in the same direction (strabismus) and a visible whiteness (leukocoria) in the normally black pupil.

Some people with familial exudative vitreoretinopathy also have reduced bone mineral density, which weakens bones and increases the risk of fractures.

### Frequency

The prevalence of familial exudative vitreoretinopathy is unknown. It appears to be rare, although affected people with normal vision may never come to medical attention.

### Causes

Mutations in the *FZD4*, *LRP5*, and *NDP* genes can cause familial exudative vitreoretinopathy. These genes provide instructions for making proteins that participate in a chemical signaling pathway that affects the way cells and tissues develop. In particular, the proteins produced from the *FZD4*, *LRP5*, and *NDP* genes appear to play critical roles in the specialization of retinal cells and the establishment of a blood supply to the retina and the inner ear. The *LRP5* protein also helps regulate bone formation.

Mutations in the *FZD4*, *LRP5*, or *NDP* gene disrupt chemical signaling during early development, which interferes with the formation of blood vessels at the edges of the retina. The resulting abnormal blood supply to this tissue leads to retinal damage and vision loss in some people with familial exudative vitreoretinopathy.

The eye abnormalities associated with familial exudative vitreoretinopathy tend to be similar no matter which gene is altered. However, affected individuals with *LRP5* gene mutations often have reduced bone mineral density in addition to vision loss. Mutations in the other genes responsible for familial exudative vitreoretinopathy do not appear to affect bone density.

In some cases, the cause of familial exudative vitreoretinopathy is unknown. Researchers believe that mutations in several as-yet-unidentified genes are responsible for the disorder in these cases.

## **Inheritance Pattern**

Familial exudative vitreoretinopathy has different inheritance patterns depending on the gene involved. Most commonly, the condition results from mutations in the *FZD4* or *LRP5* gene and has an autosomal dominant pattern of inheritance. Autosomal dominant inheritance means one copy of the altered gene in each cell is sufficient to cause the disorder. Most people with autosomal dominant familial exudative vitreoretinopathy inherit the altered gene from a parent, although the parent may not have any signs and symptoms associated with this disorder.

Familial exudative vitreoretinopathy caused by *LRP5* gene mutations can also have an autosomal recessive pattern of inheritance. Autosomal recessive inheritance means both copies of the gene in each cell have mutations. The parents of an individual with autosomal recessive familial exudative vitreoretinopathy each carry one copy of the mutated gene, but they do not have the disorder.

When familial exudative vitreoretinopathy is caused by mutations in the *NDP* gene, it has an X-linked recessive pattern of inheritance. The *NDP* gene is located on the X chromosome, which is one of the two sex chromosomes. In males (who have only one X chromosome), one altered copy of the gene in each cell is sufficient to cause the condition. In females (who have two X chromosomes), a mutation would have to occur in both copies of the gene to cause the disorder. Because it is unlikely that females will have two altered copies of this gene, males are affected by X-linked disorders much more frequently than females. A characteristic of X-linked inheritance is that fathers cannot pass X-linked traits to their sons.

## **Other Names for This Condition**

- FEVR

## **Diagnosis & Management**

### Genetic Testing Information

- What is genetic testing?  
[/primer/testing/genetictesting](#)
- Genetic Testing Registry: Exudative vitreoretinopathy 1  
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1851402/>
- Genetic Testing Registry: Exudative vitreoretinopathy 3  
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1854002/>

- Genetic Testing Registry: Exudative vitreoretinopathy 4  
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1866176/>
- Genetic Testing Registry: Familial exudative vitreoretinopathy, X-linked  
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1844579/>

#### Research Studies from ClinicalTrials.gov

- ClinicalTrials.gov  
<https://clinicaltrials.gov/ct2/results?cond=%22familial+exudative+vitreoretinopathy%22>

#### Other Diagnosis and Management Resources

- GeneReview: Familial Exudative Vitreoretinopathy, Autosomal Dominant  
<https://www.ncbi.nlm.nih.gov/books/NBK1147>
- GeneReview: NDP-Related Retinopathies  
<https://www.ncbi.nlm.nih.gov/books/NBK1331>

### **Additional Information & Resources**

#### Health Information from MedlinePlus

- Health Topic: Retinal Disorders  
<https://medlineplus.gov/retinaldisorders.html>

#### Genetic and Rare Diseases Information Center

- Familial exudative vitreoretinopathy  
<https://rarediseases.info.nih.gov/diseases/1613/familial-exudative-vitreoretinopathy>

#### Additional NIH Resources

- National Eye Institute: Diagram of the Eye  
<https://nei.nih.gov/health/eyediagram/>
- National Eye Institute: Low Vision  
<https://nei.nih.gov/health/lowvision/>

#### Educational Resources

- Orphanet: Familial exudative vitreoretinopathy  
[https://www.orpha.net/consor/cgi-bin/OC\\_Exp.php?Lng=EN&Expert=891](https://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=891)

#### Patient Support and Advocacy Resources

- American Foundation for the Blind  
<https://www.afb.org/>

### Clinical Information from GeneReviews

- Familial Exudative Vitreoretinopathy, Autosomal Dominant  
<https://www.ncbi.nlm.nih.gov/books/NBK1147>
- NDP-Related Retinopathies  
<https://www.ncbi.nlm.nih.gov/books/NBK1331>

### Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28familial+exudative+vit+reoretinopathy%5BTIAB%5D%29+OR+%28fevr%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

### Catalog of Genes and Diseases from OMIM

- EXUDATIVE VITREORETINOPATHY 1  
<http://omim.org/entry/133780>
- EXUDATIVE VITREORETINOPATHY 2, X-LINKED  
<http://omim.org/entry/305390>
- EXUDATIVE VITREORETINOPATHY 3  
<http://omim.org/entry/605750>
- EXUDATIVE VITREORETINOPATHY 4  
<http://omim.org/entry/601813>

### Medical Genetics Database from MedGen

- Familial exudative vitreoretinopathy, autosomal dominant  
<https://www.ncbi.nlm.nih.gov/medgen/468421>
- Familial exudative vitreoretinopathy, X-linked  
<https://www.ncbi.nlm.nih.gov/medgen/337030>

### **Sources for This Summary**

- Dickinson JL, Sale MM, Passmore A, FitzGerald LM, Wheatley CM, Burdon KP, Craig JE, Tengtrisor S, Carden SM, Maclean H, Mackey DA. Mutations in the NDP gene: contribution to Norrie disease, familial exudative vitreoretinopathy and retinopathy of prematurity. *Clin Experiment Ophthalmol.* 2006 Sep-Oct;34(7):682-8.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/16970763>
- Jiao X, Ventruto V, Trese MT, Shastry BS, Hejtmancik JF. Autosomal recessive familial exudative vitreoretinopathy is associated with mutations in LRP5. *Am J Hum Genet.* 2004 Nov;75(5):878-84. Epub 2004 Sep 2.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/15346351>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1182117/>

- Qin M, Hayashi H, Oshima K, Tahira T, Hayashi K, Kondo H. Complexity of the genotype-phenotype correlation in familial exudative vitreoretinopathy with mutations in the LRP5 and/or FZD4 genes. *Hum Mutat.* 2005 Aug;26(2):104-12.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/15981244>
- Robitaille J, MacDonald ML, Kaykas A, Sheldahl LC, Zeisler J, Dubé MP, Zhang LH, Singaraja RR, Guernsey DL, Zheng B, Siebert LF, Hoskin-Mott A, Trese MT, Pimstone SN, Shastry BS, Moon RT, Hayden MR, Goldberg YP, Samuels ME. Mutant frizzled-4 disrupts retinal angiogenesis in familial exudative vitreoretinopathy. *Nat Genet.* 2002 Oct;32(2):326-30. Epub 2002 Aug 12.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/12172548>
- Shukla D, Singh J, Sudheer G, Soman M, John RK, Ramasamy K, Perumalsamy N. Familial exudative vitreoretinopathy (FEVR). Clinical profile and management. *Indian J Ophthalmol.* 2003 Dec;51(4):323-8.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/14750620>
- Sims KB. NDP-Related Retinopathies. 1999 Jul 30 [updated 2014 Sep 18]. In: Pagon RA, Adam MP, Ardinger HH, Wallace SE, Amemiya A, Bean LJH, Bird TD, Ledbetter N, Mefford HC, Smith RJH, Stephens K, editors. *GeneReviews®* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2017. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1331/>  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/20301506>
- Toomes C, Bottomley HM, Jackson RM, Towns KV, Scott S, Mackey DA, Craig JE, Jiang L, Yang Z, Trembath R, Woodruff G, Gregory-Evans CY, Gregory-Evans K, Parker MJ, Black GC, Downey LM, Zhang K, Inglehearn CF. Mutations in LRP5 or FZD4 underlie the common familial exudative vitreoretinopathy locus on chromosome 11q. *Am J Hum Genet.* 2004 Apr;74(4):721-30. Epub 2004 Mar 11.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/15024691>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1181948/>
- Toomes C, Downey L. Familial Exudative Vitreoretinopathy, Autosomal Dominant. 2005 Mar 21 [updated 2011 Sep 22]. In: Pagon RA, Adam MP, Ardinger HH, Wallace SE, Amemiya A, Bean LJH, Bird TD, Ledbetter N, Mefford HC, Smith RJH, Stephens K, editors. *GeneReviews®* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2017. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1147/>  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/20301326>
- Warden SM, Andreoli CM, Mukai S. The Wnt signaling pathway in familial exudative vitreoretinopathy and Norrie disease. *Semin Ophthalmol.* 2007 Oct-Dec;22(4):211-7. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/18097984>

---

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/familial-exudative-vitreoretinopathy>

Reviewed: February 2009

Published: May 14, 2019

Lister Hill National Center for Biomedical Communications

U.S. National Library of Medicine

National Institutes of Health

Department of Health & Human Services