



## Hypomagnesemia with secondary hypocalcemia

Hypomagnesemia with secondary hypocalcemia is an inherited condition caused by the body's inability to absorb and retain magnesium that is taken in through the diet. As a result, magnesium levels in the blood are severely low (hypomagnesemia).

Hypomagnesemia impairs the function of the parathyroid glands, which are small hormone-producing glands located in the neck. Normally, the parathyroid glands release a hormone that increases blood calcium levels when they are low. Magnesium is required for the production and release of parathyroid hormone, so when magnesium is too low, insufficient parathyroid hormone is produced and blood calcium levels are also reduced (hypocalcemia). The hypocalcemia is described as "secondary" because it occurs as a consequence of hypomagnesemia.

Shortages of magnesium and calcium can cause neurological problems that begin in infancy, including painful muscle spasms (tetany) and seizures. If left untreated, hypomagnesemia with secondary hypocalcemia can lead to developmental delay, intellectual disability, a failure to gain weight and grow at the expected rate (failure to thrive), and heart failure.

### Frequency

Hypomagnesemia with secondary hypocalcemia is thought to be a rare condition, but its prevalence is unknown.

### Causes

Hypomagnesemia with secondary hypocalcemia is caused by mutations in the *TRPM6* gene. This gene provides instructions for making a protein that acts as a channel, which allows charged atoms (ions) of magnesium ( $Mg^{2+}$ ) to flow into cells; the channel may also allow small amounts of calcium ions ( $Ca^{2+}$ ) to pass into cells. Magnesium is involved in many cell processes, including production of cellular energy, maintenance of DNA building blocks (nucleotides), protein production, and cell growth and death. Magnesium and calcium are also required for the normal functioning of nerve cells that control muscle movement (motor neurons).

The TRPM6 channel is embedded in the membrane of epithelial cells that line the large intestine, structures in the kidneys known as distal convoluted tubules, the lungs, and the testes in males. When the body needs additional  $Mg^{2+}$ , the TRPM6 channel allows it to be absorbed in the intestine and filtered from the fluids that pass through the kidneys by the distal convoluted tubules. When the body has sufficient or too much  $Mg^{2+}$ , the TRPM6 channel does not filter out the  $Mg^{2+}$  from fluids but allows the ion to be released

from the kidney cells into the urine. The channel also helps to regulate  $\text{Ca}^{2+}$ , but to a lesser degree.

Most *TRPM6* gene mutations that cause hypomagnesemia with secondary hypocalcemia result in a lack of functional protein. A loss of functional TRPM6 channels prevent  $\text{Mg}^{2+}$  absorption in the intestine and cause excessive amounts of  $\text{Mg}^{2+}$  to be excreted by the kidneys and released in the urine. A lack of  $\text{Mg}^{2+}$  in the body impairs the production of parathyroid hormone, which likely reduces blood  $\text{Ca}^{2+}$  levels. Additionally, hypomagnesemia and hypocalcemia can disrupt many cell processes and impair the function of motor neurons, leading to neurological problems and movement disorders. If the condition is not effectively treated and low  $\text{Mg}^{2+}$  levels persist, signs and symptoms can worsen over time and may lead to early death.

### **Inheritance Pattern**

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

### **Other Names for This Condition**

- familial primary hypomagnesemia with hypocalcuria
- HOMG
- HSH
- hypomagnesemic tetany
- intestinal hypomagnesemia 1
- intestinal hypomagnesemia with secondary hypocalcemia

### **Diagnosis & Management**

#### Genetic Testing Information

- What is genetic testing?  
[/primer/testing/genetic-testing](#)
- Genetic Testing Registry: Hypomagnesemia 1, intestinal  
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1865974/>

#### Other Diagnosis and Management Resources

- MedlinePlus Encyclopedia: Hypomagnesemia  
<https://medlineplus.gov/ency/article/000315.htm>

## **Additional Information & Resources**

### Health Information from MedlinePlus

- Encyclopedia: Hypocalcemia--Infants  
<https://medlineplus.gov/ency/article/007229.htm>
- Encyclopedia: Hypomagnesemia  
<https://medlineplus.gov/ency/article/000315.htm>
- Health Topic: Fluid and Electrolyte Balance  
<https://medlineplus.gov/fluidandelectrolytebalance.html>

### Additional NIH Resources

- National Institute of Neurological Disorders and Stroke: Epilepsy Information Page  
<https://www.ninds.nih.gov/Disorders/All-Disorders/Epilepsy-Information-Page>

### Educational Resources

- Boston Children's Hospital: Seizures in Children  
<http://www.childrenshospital.org/conditions-and-treatments/conditions/s/seizures>
- Centers for Disease Control and Prevention: Intellectual Disability  
[https://www.cdc.gov/ncbddd/actearly/pdf/parents\\_pdfs/IntellectualDisability.pdf](https://www.cdc.gov/ncbddd/actearly/pdf/parents_pdfs/IntellectualDisability.pdf)
- Cincinnati Children's Hospital: Epilepsy and Seizures  
<https://www.cincinnatichildrens.org/health/e/epilepsy-seizures>
- Johns Hopkins Medicine: Hypoparathyroidism  
<https://www.hopkinsmedicine.org/health/conditions-and-diseases/hypoparathyroidism>
- Kennedy Krieger Institute: Epilepsy (Seizure Disorder)  
<https://www.kennedykrieger.org/patient-care/conditions/epilepsy-seizure-disorder>
- Kennedy Krieger Institute: Intellectual Disability  
<https://www.kennedykrieger.org/patient-care/conditions/intellectual-disability>
- MalaCards: hypomagnesemia 1, intestinal  
[https://www.malacards.org/card/hypomagnesemia\\_1\\_intestinal](https://www.malacards.org/card/hypomagnesemia_1_intestinal)
- Merck Manual Consumer Version: Hypocalcemia  
<https://www.merckmanuals.com/home/hormonal-and-metabolic-disorders/electrolyte-balance/hypocalcemia-low-level-of-calcium-in-the-blood>
- Merck Manual Professional Version: Hypomagnesemia  
<https://www.merckmanuals.com/professional/endocrine-and-metabolic-disorders/electrolyte-disorders/hypomagnesemia>
- Orphanet: Primary hypomagnesemia with secondary hypocalcemia  
[https://www.orpha.net/consor/cgi-bin/OC\\_Exp.php?Lng=EN&Expert=30924](https://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=30924)

## Patient Support and Advocacy Resources

- American Association on Intellectual and Developmental Disabilities (AAIDD)  
<http://aaid.org/>
- American Epilepsy Society  
<https://www.aesnet.org/>
- CURE Epilepsy  
<https://www.cureepilepsy.org/>

## Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28hypomagnesemia+with+secondary+hypocalcemia%5BTIAB%5D%29+OR+%28intestinal+hypomagnesemia%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

## Catalog of Genes and Diseases from OMIM

- HYPOMAGNESEMIA 1, INTESTINAL  
<http://omim.org/entry/602014>

## **Sources for This Summary**

- Chubanov V, Gudermann T, Schlingmann KP. Essential role for TRPM6 in epithelial magnesium transport and body magnesium homeostasis. *Pflugers Arch*. 2005 Oct;451(1):228-34. Epub 2005 Jun 17. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/16075242>
- Katayama K, Povalko N, Yatsuga S, Nishioka J, Kakuma T, Matsuishi T, Koga Y. New TRPM6 mutation and management of hypomagnesaemia with secondary hypocalcaemia. *Brain Dev*. 2015 Mar;37(3):292-8. doi: 10.1016/j.braindev.2014.06.006. Epub 2014 Jun 28.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/24985022>
- Lainez S, Schlingmann KP, van der Wijst J, Dworniczak B, van Zeeland F, Konrad M, Bindels RJ, Hoenderop JG. New TRPM6 missense mutations linked to hypomagnesemia with secondary hypocalcemia. *Eur J Hum Genet*. 2014 Apr;22(4):497-504. doi: 10.1038/ejhg.2013.178. Epub 2013 Aug 14.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/23942199>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3953905/>
- Schlingmann KP, Sassen MC, Weber S, Pechmann U, Kusch K, Pelken L, Lotan D, Syrrou M, Prebble JJ, Cole DE, Metzger DL, Rahman S, Tajima T, Shu SG, Waldegger S, Seyberth HW, Konrad M. Novel TRPM6 mutations in 21 families with primary hypomagnesemia and secondary hypocalcemia. *J Am Soc Nephrol*. 2005 Oct;16(10):3061-9. Epub 2005 Aug 17.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/16107578>
- Schlingmann KP, Weber S, Peters M, Niemann Nejsum L, Vitzthum H, Klingel K, Kratz M, Haddad E, Ristoff E, Dinour D, Syrrou M, Nielsen S, Sassen M, Waldegger S, Seyberth HW, Konrad M. Hypomagnesemia with secondary hypocalcemia is caused by mutations in TRPM6, a new member of the TRPM gene family. *Nat Genet*. 2002 Jun;31(2):166-70. Epub 2002 May 28.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/12032568>

---

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/hypomagnesemia-with-secondary-hypocalcemia>

Reviewed: January 2015

Published: September 10, 2019

Lister Hill National Center for Biomedical Communications

U.S. National Library of Medicine

National Institutes of Health

Department of Health & Human Services