



Maturity-onset diabetes of the young

Maturity-onset diabetes of the young (MODY) is a group of several conditions characterized by abnormally high blood sugar levels. These forms of diabetes typically begin before age 30, although they can occur later in life. In MODY, elevated blood sugar arises from reduced production of insulin, which is a hormone produced in the pancreas that helps regulate blood sugar levels. Specifically, insulin controls how much glucose (a type of sugar) is passed from the blood into cells, where it is used as an energy source.

The different types of MODY are distinguished by their genetic causes. The most common types are *HNF1A*-MODY (also known as MODY3), accounting for 50 to 70 percent of cases, and *GCK*-MODY (MODY2), accounting for 30 to 50 percent of cases. Less frequent types include *HNF4A*-MODY (MODY1) and renal cysts and diabetes (RCAD) syndrome (also known as *HNF1B*-MODY or MODY5), which each account for 5 to 10 percent of cases. At least ten other types have been identified, and these are very rare.

HNF1A-MODY and *HNF4A*-MODY have similar signs and symptoms that develop slowly over time. Early signs and symptoms in these types are caused by high blood sugar and may include frequent urination (polyuria), excessive thirst (polydipsia), fatigue, blurred vision, weight loss, and recurrent skin infections. Over time uncontrolled high blood sugar can damage small blood vessels in the eyes and kidneys. Damage to the light-sensitive tissue at the back of the eye (the retina) causes a condition known as diabetic retinopathy that can lead to vision loss and eventual blindness. Kidney damage (diabetic nephropathy) can lead to kidney failure and end-stage renal disease (ESRD). While these two types of MODY are very similar, certain features are particular to each type. For example, babies with *HNF4A*-MODY tend to weigh more than average or have abnormally low blood sugar at birth, even though other signs of the condition do not occur until childhood or young adulthood. People with *HNF1A*-MODY have a higher-than-average risk of developing noncancerous (benign) liver tumors known as hepatocellular adenomas.

GCK-MODY is a very mild type of the condition. People with this type have slightly elevated blood sugar levels, particularly in the morning before eating (fasting blood sugar). However, affected individuals often have no symptoms related to the disorder, and diabetes-related complications are extremely rare.

RCAD is associated with a combination of diabetes and kidney or urinary tract abnormalities (unrelated to the elevated blood sugar), most commonly fluid-filled sacs (cysts) in the kidneys. However, the signs and symptoms are variable, even within families, and not everyone with RCAD has both features. Affected individuals may have

other features unrelated to diabetes, such as abnormalities of the pancreas or liver or a form of arthritis called gout.

Frequency

MODY is estimated to account for 1 to 3 percent of all cases of diabetes.

Causes

MODY can be caused by a mutation in one of several genes. *HNF1A*-MODY, *GCK*-MODY, *HNF4A*-MODY, and RCAD, are caused by mutations in the *HNF1A*, *GCK*, *HNF4A*, and *HNF1B* gene, respectively. All of these genes provide instructions for making proteins involved in the production of insulin to control blood sugar levels in the body. In particular, the proteins are important in specialized cells in the pancreas called beta cells, which secrete insulin.

The proteins produced from the *HNF1A*, *HNF4A*, and *HNF1B* genes all act as transcription factors, which means they control the activity of other genes. In particular, these proteins regulate genes that direct the development and function of beta cells. *HNF1A*, *HNF4A*, or *HNF1B* gene mutations result in production of an altered transcription factor that is unable to function normally. These changes alter gene activity in cells, impairing normal beta cell development and function. As a result, beta cells are less able than normal to produce insulin in response to sugar in the blood, which means the body cannot control blood sugar. Elevated blood sugar results in the signs and symptoms of MODY. Some of these MODY-related genes play roles in the development of other body systems, in addition to beta cells. Disrupted development of these systems underlies additional signs and symptoms in particular forms of MODY. For example, the *HNF1B* gene is involved in kidney development, which helps explain the kidney abnormalities in people with RCAD.

The protein produced from the *GCK* gene acts as a sensor that recognizes when the amount of glucose in the blood rises. In response, the protein helps stimulate the release of insulin from beta cells so sugar can be taken up and used by cells for energy. This protein also helps determine when excess sugar should be taken into liver cells and stored. Mutations in the *GCK* gene limit the protein's ability to sense a rise in blood sugar, so levels remain elevated.

Other genes involved in controlling blood sugar cause rare types of MODY. It is likely that additional genes that have not been identified are also involved in the condition.

Inheritance Pattern

MODY is inherited in an autosomal dominant pattern, which means one copy of the altered gene in each cell is sufficient to cause the disorder.

In most cases, an affected person inherits the mutation from one affected parent. Other cases result from new mutations in the gene and occur in people with no history of the disorder in their family.

Other Names for This Condition

- MODY

Diagnosis & Management

Genetic Testing Information

- What is genetic testing?
[/primer/testing/genetictesting](#)
- Genetic Testing Registry: Maturity onset diabetes mellitus in young
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0342276/>
- Genetic Testing Registry: Maturity-onset diabetes of the young, type 1
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1852093/>
- Genetic Testing Registry: Maturity-onset diabetes of the young, type 2
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1841962/>
- Genetic Testing Registry: Maturity-onset diabetes of the young, type 3
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1838100/>

Research Studies from ClinicalTrials.gov

- ClinicalTrials.gov
<https://clinicaltrials.gov/ct2/results?cond=%22maturity-onset+diabetes+of+the+young%22>

Other Diagnosis and Management Resources

- GeneReview: Maturity-Onset Diabetes of the Young Overview
<https://www.ncbi.nlm.nih.gov/books/NBK500456>

Additional Information & Resources

Health Information from MedlinePlus

- Health Topic: Blood Sugar
<https://medlineplus.gov/bloodsugar.html>
- Health Topic: Diabetes
<https://medlineplus.gov/diabetes.html>
- Medical Tests: Blood Glucose Test
<https://medlineplus.gov/lab-tests/blood-glucose-test/>
- Medical Tests: Insulin in Blood
<https://medlineplus.gov/lab-tests/insulin-in-blood/>

Genetic and Rare Diseases Information Center

- Maturity-onset diabetes of the young
<https://rarediseases.info.nih.gov/diseases/3697/maturity-onset-diabetes-of-the-young>

Additional NIH Resources

- National Institute of Diabetes and Digestive and Kidney Diseases: Monogenic Diabetes (Neonatal Diabetes Mellitus & MODY)
<https://www.niddk.nih.gov/health-information/diabetes/overview/what-is-diabetes/monogenic-neonatal-mellitus-mody>

Educational Resources

- MalaCards: maturity-onset diabetes of the young
https://www.malacards.org/card/maturity_onset_diabetes_of_the_young

Patient Support and Advocacy Resources

- American Diabetes Association
<https://www.diabetes.org/>
- Diabetes UK
<https://www.diabetes.org.uk/>
- Kovler Diabetes Center at the University of Chicago: Monogenic Diabetes Registry
<https://kovlerdiabetescenter.org/for-patients/individuals-with-monogenic-diabetes/>

Clinical Information from GeneReviews

- Maturity-Onset Diabetes of the Young Overview
<https://www.ncbi.nlm.nih.gov/books/NBK500456>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28Diabetes+Mellitus%5BMAJR%5D%29+AND+%28maturity-onset+diabetes+of+the+young%5BTIAB%5D%29+AND+english%5BIa%5D+AND+human%5Bmh%5D+AND+%22last+720+days%22%5Bdp%5D>

Catalog of Genes and Diseases from OMIM

- MATURITY-ONSET DIABETES OF THE YOUNG
<http://omim.org/entry/606391>
- MATURITY-ONSET DIABETES OF THE YOUNG, TYPE 1
<http://omim.org/entry/125850>
- MATURITY-ONSET DIABETES OF THE YOUNG, TYPE 2
<http://omim.org/entry/125851>

- MATURITY-ONSET DIABETES OF THE YOUNG, TYPE 3
<http://omim.org/entry/600496>
- RENAL CYSTS AND DIABETES SYNDROME
<http://omim.org/entry/137920>

Medical Genetics Database from MedGen

- Maturity onset diabetes mellitus in young
<https://www.ncbi.nlm.nih.gov/medgen/87433>

Sources for This Summary

- Froguel P, Zouali H, Vionnet N, Velho G, Vaxillaire M, Sun F, Lesage S, Stoffel M, Takeda J, Passa P, et al. Familial hyperglycemia due to mutations in glucokinase. Definition of a subtype of diabetes mellitus. *N Engl J Med.* 1993 Mar 11;328(10):697-702.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/8433729>
- Haliyur R, Tong X, Sanyoura M, Shrestha S, Lindner J, Saunders DC, Aramandla R, Poffenberger G, Redick SD, Bottino R, Prasad N, Levy SE, Blind RD, Harlan DM, Philipson LH, Stein RW, Brissova M, Powers AC. Human islets expressing HNF1A variant have defective β cell transcriptional regulatory networks. *J Clin Invest.* 2019 Jan 2;129(1):246-251. doi: 10.1172/JCI121994. Epub 2018 Dec 3.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/30507613>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6307934/>
- Malikova J, Kaci A, Dusatkova P, Aukrust I, Torsvik J, Vesela K, Kankova PD, Njølstad PR, Pruhova S, Bjørkhaug L. Functional Analyses of HNF1A-MODY Variants Refine the Interpretation of Identified Sequence Variants. *J Clin Endocrinol Metab.* 2020 Apr 1;105(4). pii: dgaa051. doi: 10.1210/clinem/dgaa051.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/32017842>
- Naylor R, Knight Johnson A, del Gaudio D. Maturity-Onset Diabetes of the Young Overview. 2018 May 24. In: Adam MP, Ardinger HH, Pagon RA, Wallace SE, Bean LJH, Stephens K, Amemiya A, editors. *GeneReviews®* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2020. Available from <http://www.ncbi.nlm.nih.gov/books/NBK500456/>
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/29792621>
- Negahdar M, Aukrust I, Molnes J, Solheim MH, Johansson BB, Sagen JV, Dahl-Jørgensen K, Kulkarni RN, Søvik O, Flatmark T, Njølstad PR, Bjørkhaug L. GCK-MODY diabetes as a protein misfolding disease: the mutation R275C promotes protein misfolding, self-association and cellular degradation. *Mol Cell Endocrinol.* 2014 Jan 25;382(1):55-65. doi: 10.1016/j.mce.2013.08.020. Epub 2013 Aug 31.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/24001579>
- Osbak KK, Colclough K, Saint-Martin C, Beer NL, Bellanné-Chantelot C, Ellard S, Gloyn AL. Update on mutations in glucokinase (GCK), which cause maturity-onset diabetes of the young, permanent neonatal diabetes, and hyperinsulinemic hypoglycemia. *Hum Mutat.* 2009 Nov;30(11):1512-26. doi: 10.1002/humu.21110. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19790256>
- Singh P, Tung SP, Han EH, Lee IK, Chi YI. Dimerization defective MODY mutations of hepatocyte nuclear factor 4 α . *Mutat Res.* 2019 Mar;814:1-6. doi: 10.1016/j.mrfmmm.2019.01.002. Epub 2019 Jan 9.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/30648609>

- Yahaya TO, Ufuoma SB. Genetics and Pathophysiology of Maturity-onset Diabetes of the Young (MODY): A Review of Current Trends. Oman Med J. 2020 May 28;35(3):e126. doi: 10.5001/omj.2020.44. eCollection 2020 May. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/32489678>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7254248/>
 - Yamagata K. Roles of HNF1 α and HNF4 α in pancreatic β -cells: lessons from a monogenic form of diabetes (MODY). Vitam Horm. 2014;95:407-23. doi: 10.1016/B978-0-12-800174-5.00016-8. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/24559927>
-

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/maturity-onset-diabetes-of-the-young>

Reviewed: July 2020

Published: August 17, 2020

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services