



Prothrombin thrombophilia

Prothrombin thrombophilia is an inherited disorder of blood clotting. Thrombophilia is an increased tendency to form abnormal blood clots in blood vessels. People who have prothrombin thrombophilia are at somewhat higher than average risk for a type of clot called a deep venous thrombosis, which typically occurs in the deep veins of the legs. Affected people also have an increased risk of developing a pulmonary embolism, which is a clot that travels through the bloodstream and lodges in the lungs. Most people with prothrombin thrombophilia never develop abnormal blood clots, however.

Some research suggests that prothrombin thrombophilia is associated with a somewhat increased risk of pregnancy loss (miscarriage) and may also increase the risk of other complications during pregnancy. These complications may include pregnancy-induced high blood pressure (preeclampsia), slow fetal growth, and early separation of the placenta from the uterine wall (placental abruption). It is important to note, however, that most women with prothrombin thrombophilia have normal pregnancies.

Frequency

Prothrombin thrombophilia is the second most common inherited form of thrombophilia after factor V Leiden thrombophilia. Approximately 1 in 50 people in the white population in the United States and Europe has prothrombin thrombophilia. This condition is less common in other ethnic groups, occurring in less than one percent of African American, Native American, or Asian populations.

Causes

Prothrombin thrombophilia is caused by a particular mutation in the *F2* gene. The *F2* gene plays a critical role in the formation of blood clots in response to injury. The protein produced from the *F2* gene, prothrombin (also called coagulation factor II), is the precursor to a protein called thrombin that initiates a series of chemical reactions in order to form a blood clot. The particular mutation that causes prothrombin thrombophilia results in an overactive *F2* gene that causes too much prothrombin to be produced. An abundance of prothrombin leads to more thrombin, which promotes the formation of blood clots.

Other factors also increase the risk of blood clots in people with prothrombin thrombophilia. These factors include increasing age, obesity, trauma, surgery, smoking, the use of oral contraceptives (birth control pills) or hormone replacement therapy, and pregnancy. The combination of prothrombin thrombophilia and mutations in other genes involved in blood clotting can also influence risk.

Inheritance Pattern

The risk of developing an abnormal clot in a blood vessel depends on whether a person inherits one or two copies of the *F2* gene mutation that causes prothrombin thrombophilia. In the general population, the risk of developing an abnormal blood clot is about 1 in 1,000 people per year. Inheriting one copy of the *F2* gene mutation increases that risk to 2 to 3 in 1,000. People who inherit two copies of the mutation, one from each parent, may have a risk as high as 20 in 1,000.

Other Names for This Condition

- hyperprothrombinemia
- Prothrombin G20210A Thrombophilia

Diagnosis & Management

Genetic Testing Information

- What is genetic testing?
[/primer/testing/genetictesting](#)
- Genetic Testing Registry: Hereditary factor II deficiency disease
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0272317/>

Research Studies from ClinicalTrials.gov

- ClinicalTrials.gov
<https://clinicaltrials.gov/ct2/results?cond=%22Thrombophilia%22+OR+%22prothrombin+thrombophilia%22>

Other Diagnosis and Management Resources

- GeneReview: Prothrombin-Related Thrombophilia
<https://www.ncbi.nlm.nih.gov/books/NBK1148>
- MedlinePlus Encyclopedia: Deep venous thrombosis
<https://medlineplus.gov/ency/article/000156.htm>
- MedlinePlus Encyclopedia: Pulmonary embolus
<https://medlineplus.gov/ency/article/000132.htm>

Additional Information & Resources

Health Information from MedlinePlus

- Encyclopedia: Deep venous thrombosis
<https://medlineplus.gov/ency/article/000156.htm>
- Encyclopedia: Pulmonary embolus
<https://medlineplus.gov/ency/article/000132.htm>
- Health Topic: Blood Clots
<https://medlineplus.gov/bloodclots.html>

Genetic and Rare Diseases Information Center

- Prothrombin-related thrombophilia
<https://rarediseases.info.nih.gov/diseases/10815/prothrombin-related-thrombophilia>

Additional NIH Resources

- National Heart, Lung, and Blood Institute: Deep Vein Thrombosis
<https://www.nhlbi.nih.gov/health-topics/venous-thromboembolism>

Educational Resources

- MalaCards: prothrombin-related thrombophilia
https://www.malacards.org/card/prothrombin_related_thrombophilia
- March of Dimes: Thrombophilias
<https://www.marchofdimes.org/pregnancy/thrombophilias.aspx>
- Merck Manual for Patients and Caregivers: Thrombophilia
<https://www.merckmanuals.com/home/blood-disorders/excessive-clotting/excessive-clotting>
- Orphanet: Congenital factor II deficiency
https://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=325
- Prothrombin Cardiology Patient Page
<https://www.ahajournals.org/doi/full/10.1161/01.cir.0000135582.53444.87>

Patient Support and Advocacy Resources

- National Blood Clot Alliance
<https://www.stoptheclot.org/>

Clinical Information from GeneReviews

- Prothrombin-Related Thrombophilia
<https://www.ncbi.nlm.nih.gov/books/NBK1148>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28prothrombin%5BTIAB%5D%29+AND+%28thrombophilia%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

Catalog of Genes and Diseases from OMIM

- COAGULATION FACTOR II
<http://omim.org/entry/176930>
- THROMBOPHILIA DUE TO THROMBIN DEFECT
<http://omim.org/entry/188050>

Medical Genetics Database from MedGen

- Hereditary factor II deficiency disease
<https://www.ncbi.nlm.nih.gov/medgen/124425>

Sources for This Summary

- Bosler D, Mattson J, Crisan D. Phenotypic Heterogeneity in Patients with Homozygous Prothrombin 20210AA Genotype. A paper from the 2005 William Beaumont Hospital Symposium on Molecular Pathology. J Mol Diagn. 2006 Sep;8(4):420-5. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16931580>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1867627/>
- McGlennen RC, Key NS. Clinical and laboratory management of the prothrombin G20210A mutation. Arch Pathol Lab Med. 2002 Nov;126(11):1319-25. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12421139>
- Spector EB, Grody WW, Matteson CJ, Palomaki GE, Bellissimo DB, Wolff DJ, Bradley LA, Prior TW, Feldman G, Popovich BW, Watson MS, Richards CS. Technical standards and guidelines: venous thromboembolism (Factor V Leiden and prothrombin 20210G >A testing): a disease-specific supplement to the standards and guidelines for clinical genetics laboratories. Genet Med. 2005 Jul-Aug;7(6):444-53.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16024978>
- Varga EA, Moll S. Cardiology patient pages. Prothrombin 20210 mutation (factor II mutation). Circulation. 2004 Jul 20;110(3):e15-8.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15262854>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/prothrombin-thrombophilia>

Reviewed: August 2008

Published: May 14, 2019

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services