



## Tyrosinemia

Tyrosinemia is a genetic disorder characterized by disruptions in the multistep process that breaks down the amino acid tyrosine, a building block of most proteins. If untreated, tyrosine and its byproducts build up in tissues and organs, which can lead to serious health problems.

There are three types of tyrosinemia, which are each distinguished by their symptoms and genetic cause. Tyrosinemia type I, the most severe form of this disorder, is characterized by signs and symptoms that begin in the first few months of life. Affected infants fail to gain weight and grow at the expected rate (failure to thrive) due to poor food tolerance because high-protein foods lead to diarrhea and vomiting. Affected infants may also have yellowing of the skin and whites of the eyes (jaundice), a cabbage-like odor, and an increased tendency to bleed (particularly nosebleeds). Tyrosinemia type I can lead to liver and kidney failure, softening and weakening of the bones (rickets), and an increased risk of liver cancer (hepatocellular carcinoma). Some affected children have repeated neurologic crises that consist of changes in mental state, reduced sensation in the arms and legs (peripheral neuropathy), abdominal pain, and respiratory failure. These crises can last from 1 to 7 days. Untreated, children with tyrosinemia type I often do not survive past the age of 10.

Tyrosinemia type II can affect the eyes, skin, and mental development. Signs and symptoms often begin in early childhood and include eye pain and redness, excessive tearing, abnormal sensitivity to light (photophobia), and thick, painful skin on the palms of their hands and soles of their feet (palmoplantar hyperkeratosis). About 50 percent of individuals with tyrosinemia type II have some degree of intellectual disability.

Tyrosinemia type III is the rarest of the three types. The characteristic features of this type include intellectual disability, seizures, and periodic loss of balance and coordination (intermittent ataxia).

About 10 percent of newborns have temporarily elevated levels of tyrosine (transient tyrosinemia). In these cases, the cause is not genetic. The most likely causes are vitamin C deficiency or immature liver enzymes due to premature birth.

### Frequency

Worldwide, tyrosinemia type I affects about 1 in 100,000 individuals. This type is more common in Norway where 1 in 60,000 to 74,000 individuals are affected. Tyrosinemia type I is even more common in Quebec, Canada where it occurs in about 1 in 16,000 individuals. In the Saguenay-Lac St. Jean region of Quebec, tyrosinemia type I affects 1 in 1,846 people.

Tyrosinemia type II occurs in fewer than 1 in 250,000 individuals worldwide. Tyrosinemia type III is very rare; only a few cases have been reported.

## Genetic Changes

Mutations in the *FAH*, *TAT*, and *HPD* genes can cause tyrosinemia types I, II, and III, respectively.

In the liver, enzymes break down tyrosine in a five step process, resulting in molecules that are either excreted by the kidneys or used to produce energy or make other substances in the body. The *FAH* gene provides instructions for the fumarylacetoacetate hydrolase enzyme, which is responsible for the final step of tyrosine breakdown. The enzyme produced from the *TAT* gene, called tyrosine aminotransferase enzyme, is involved at the first step in the process. The *HPD* gene provides instructions for making the 4-hydroxyphenylpyruvate dioxygenase enzyme, which is responsible for the second step.

Mutations in the *FAH*, *TAT*, or *HPD* gene cause a decrease in the activity of one of the enzymes in the breakdown of tyrosine. As a result, tyrosine and its byproducts accumulate to toxic levels, which can cause damage and death to cells in the liver, kidneys, nervous system, and other organs.

## Inheritance Pattern

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

## Other Names for This Condition

- hereditary tyrosinemia
- hypertyrosinaemia
- hypertyrosinemia
- tyrosinaemia

## Diagnosis & Management

### Formal Diagnostic Criteria

- ACT Sheet: Increased Tyrosine  
<https://www.ncbi.nlm.nih.gov/books/NBK55827/bin/Tyrosine.pdf>

## Genetic Testing

- Genetic Testing Registry: 4-Hydroxyphenylpyruvate dioxygenase deficiency  
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0268623/>
- Genetic Testing Registry: Tyrosinemia type 2  
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0268487/>
- Genetic Testing Registry: Tyrosinemia type I  
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0268490/>

## Other Diagnosis and Management Resources

- Baby's First Test: Tyrosinemia, Type I  
<http://www.babysfirsttest.org/newborn-screening/conditions/tyrosinemia-type-i>
- Baby's First Test: Tyrosinemia, Type II  
<http://www.babysfirsttest.org/newborn-screening/conditions/tyrosinemia-type-ii>
- Baby's First Test: Tyrosinemia, Type III  
<http://www.babysfirsttest.org/newborn-screening/conditions/tyrosinemia-type-iii>
- GeneReview: Tyrosinemia Type I  
<https://www.ncbi.nlm.nih.gov/books/NBK1515>
- MedlinePlus Encyclopedia: Aminoaciduria  
<https://medlineplus.gov/ency/article/003366.htm>

## General Information from MedlinePlus

- Diagnostic Tests  
<https://medlineplus.gov/diagnostictests.html>
- Drug Therapy  
<https://medlineplus.gov/drugtherapy.html>
- Genetic Counseling  
<https://medlineplus.gov/geneticcounseling.html>
- Palliative Care  
<https://medlineplus.gov/palliativecare.html>
- Surgery and Rehabilitation  
<https://medlineplus.gov/surgeryandrehabilitation.html>

## **Additional Information & Resources**

### MedlinePlus

- Encyclopedia: Aminoaciduria  
<https://medlineplus.gov/ency/article/003366.htm>
- Health Topic: Amino Acid Metabolism Disorders  
<https://medlineplus.gov/aminoacidmetabolismdisorders.html>

- Health Topic: Liver Diseases  
<https://medlineplus.gov/liverdiseases.html>
- Health Topic: Newborn Screening  
<https://medlineplus.gov/newbornscreening.html>

### Genetic and Rare Diseases Information Center

- Tyrosinemia type 1  
<https://rarediseases.info.nih.gov/diseases/2658/tyrosinemia-type-1>
- Tyrosinemia type 2  
<https://rarediseases.info.nih.gov/diseases/3105/tyrosinemia-type-2>
- Tyrosinemia type 3  
<https://rarediseases.info.nih.gov/diseases/10332/tyrosinemia-type-3>

### Educational Resources

- CLIMB: Tyrosinaemia Type 1 Info Sheet  
<http://www.climb.org.uk/wp-content/uploads/2017/09/Tyrosinaemia-Type-1.pdf>
- Disease InfoSearch: Tyrosinemia type 1  
<http://www.diseaseinfosearch.org/Tyrosinemia+type+1/7281>
- Disease InfoSearch: Tyrosinemia type 2  
<http://www.diseaseinfosearch.org/Tyrosinemia+type+2/7282>
- Disease InfoSearch: Tyrosinemia type 3  
<http://www.diseaseinfosearch.org/Tyrosinemia+type+3/7283>
- Illinois Department of Public Health  
<http://www.idph.state.il.us/HealthWellness/fs/tyrosinemia.htm>
- MalaCards: tyrosinemia  
<http://www.malacards.org/card/tyrosinemia>
- MalaCards: tyrosinemia, type i  
[http://www.malacards.org/card/tyrosinemia\\_type\\_i\\_2](http://www.malacards.org/card/tyrosinemia_type_i_2)
- MalaCards: tyrosinemia, type ii  
[http://www.malacards.org/card/tyrosinemia\\_type\\_ii\\_2](http://www.malacards.org/card/tyrosinemia_type_ii_2)
- MalaCards: tyrosinemia, type iii  
[http://www.malacards.org/card/tyrosinemia\\_type\\_iii\\_2](http://www.malacards.org/card/tyrosinemia_type_iii_2)
- Orphanet: Tyrosinemia type 1  
[https://www.orpha.net/consor/cgi-bin/OC\\_Exp.php?Lng=EN&Expert=882](https://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=882)
- Orphanet: Tyrosinemia type 2  
[https://www.orpha.net/consor/cgi-bin/OC\\_Exp.php?Lng=EN&Expert=28378](https://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=28378)
- Screening, Technology, and Research in Genetics  
<http://www.newbornscreening.info/Parents/aminoaciddisorders/Tyrosinemia.html>

- The Swedish Information Centre for Rare Diseases  
<http://www.socialstyrelsen.se/rarediseases/tyrosinemiatype1>
- University of Washington  
<http://depts.washington.edu/tyros/abouttyr.htm>
- Virginia Department of Health  
[http://www.vdh.virginia.gov/content/uploads/sites/33/2016/11/Parent-Fact-Sheet\\_TYR-I\\_English.pdf](http://www.vdh.virginia.gov/content/uploads/sites/33/2016/11/Parent-Fact-Sheet_TYR-I_English.pdf)

#### Patient Support and Advocacy Resources

- Canadian Liver Foundation  
<https://www.liver.ca/patients-caregivers/liver-diseases/tyrosinemia/>
- CLIMB: Children Living with Inherited Metabolic Diseases (UK)  
<http://www.climb.org.uk>
- National Organization for Rare Disorders (NORD)  
<https://rarediseases.org/rare-diseases/tyrosinemia-type-1/>
- University of Washington: Tyrosinemia Pal  
<http://depts.washington.edu/tyros/tyrpal.htm>

#### GeneReviews

- Tyrosinemia Type I  
<https://www.ncbi.nlm.nih.gov/books/NBK1515>

#### ClinicalTrials.gov

- ClinicalTrials.gov  
<https://clinicaltrials.gov/ct2/results?cond=%22tyrosinemia%22>

#### Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28Tyrosinemias%5BMAJR%5D%29+AND+%28tyrosinemia%5BTIAB%5D%29+AND+english%5BIa%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

#### OMIM

- TYROSINEMIA, TYPE I  
<http://omim.org/entry/276700>
- TYROSINEMIA, TYPE II  
<http://omim.org/entry/276600>
- TYROSINEMIA, TYPE III  
<http://omim.org/entry/276710>

## Sources for This Summary

- Bouyacoub Y, Zribi H, Azzouz H, Nasrallah F, Abdelaziz RB, Kacem M, Rekaya B, Messaoud O, Romdhane L, Charfeddine C, Bouziri M, Bouziri S, Tebib N, Mokni M, Kaabachi N, Boubaker S, Abdelhak S. Novel and recurrent mutations in the TAT gene in Tunisian families affected with Richner-Hanhart syndrome. *Gene*. 2013 Oct 15;529(1):45-9. doi: 10.1016/j.gene.2013.07.066. Epub 2013 Aug 13.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/23954227>
- Couce ML, Dalmau J, del Toro M, Pintos-Morell G, Aldámiz-Echevarría L; Spanish Working Group on Tyrosinemia type 1. Tyrosinemia type 1 in Spain: mutational analysis, treatment and long-term outcome. *Pediatr Int*. 2011 Dec;53(6):985-9. doi: 10.1111/j.1442-200X.2011.03427.x.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/21752152>
- Fernández-Lainez C, Ibarra-González I, Belmont-Martínez L, Monroy-Santoyo S, Guillén-López S, Vela-Amieva M. Tyrosinemia type I: clinical and biochemical analysis of patients in Mexico. *Ann Hepatol*. 2014 Mar-Apr;13(2):265-72.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/24552869>
- Heylen E, Scherer G, Vincent MF, Marie S, Fischer J, Nassogne MC. Tyrosinemia Type III detected via neonatal screening: management and outcome. *Mol Genet Metab*. 2012 Nov;107(3):605-7. doi: 10.1016/j.ymgme.2012.09.002. Epub 2012 Sep 7.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/23036342>
- Macsai MS, Schwartz TL, Hinkle D, Hummel MB, Mulhern MG, Rootman D. Tyrosinemia type II: nine cases of ocular signs and symptoms. *Am J Ophthalmol*. 2001 Oct;132(4):522-7.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/11589874>
- de Laet C, Dionisi-Vici C, Leonard JV, McKiernan P, Mitchell G, Monti L, de Baulny HO, Pintos-Morell G, Spiekorkötter U. Recommendations for the management of tyrosinaemia type 1. *Orphanet J Rare Dis*. 2013 Jan 11;8:8. doi: 10.1186/1750-1172-8-8. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/23311542>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3558375/>

---

Reprinted from Genetics Home Reference:  
<https://ghr.nlm.nih.gov/condition/tyrosinemia>

Reviewed: August 2015  
Published: July 17, 2018

Lister Hill National Center for Biomedical Communications  
U.S. National Library of Medicine  
National Institutes of Health  
Department of Health & Human Services