



ABCC9 gene

ATP binding cassette subfamily C member 9

Normal Function

The *ABCC9* gene provides instructions for making the sulfonylurea receptor 2 (SUR2) protein. This protein forms one part (subunit) of a channel that transports charged atoms of potassium (potassium ions) across cell membranes. Each of these channels consists of eight subunits: four SUR2 proteins and four proteins produced from either the *KCNJ8* or *KCNJ11* gene. The SUR2 subunits regulate the activity of the channel, determining whether it is open or closed.

Channels made with the SUR2 protein are known as ATP-sensitive potassium (K-ATP) channels. The channels open and close in response to the amount of ATP, the cell's main energy source, inside the cell. The resulting transport of potassium ions is part of a complex network of signals that relay chemical messages into and out of cells.

Although K-ATP channels are present in cells and tissues throughout the body, the highest levels of SUR2-containing channels are found in skeletal and heart (cardiac) muscle. These channels indirectly help regulate the concentration of calcium ions in cells. This regulation is essential for normal heart function. The function of these channels in other tissues is unclear.

Health Conditions Related to Genetic Changes

Cantú syndrome

At least 14 mutations in the *ABCC9* gene have been found to cause Cantú syndrome, a rare condition characterized by excess hair growth (hypertrichosis), a distinctive facial appearance, and heart defects. Each of the mutations changes a single protein building block (amino acid) in the SUR2 protein. These changes likely alter the structure of the protein and its ability to regulate the activity of K-ATP channels. Studies suggest that the abnormal channels are open when they should be closed. However, it is unknown how this problem with potassium channel function leads to excess hair growth, heart defects, and the other features of Cantú syndrome.

familial dilated cardiomyopathy

other disorders


At least two mutations in the *ABCC9* gene have been identified in people with dilated cardiomyopathy, a form of heart disease that enlarges and weakens the cardiac muscle, preventing the heart from pumping blood efficiently. Signs and symptoms of

this condition can include an irregular heartbeat (arrhythmia), shortness of breath, extreme tiredness (fatigue), and swelling of the legs and feet. Research suggests that each mutation changes the structure of the SUR2 protein and disrupts the regulation of the K-ATP channel. Although K-ATP channels appear to play an important role in cardiac muscle, little is known about how malfunctioning channels are related to dilated cardiomyopathy.

Chromosomal Location

Cytogenetic Location: 12p12.1, which is the short (p) arm of chromosome 12 at position 12.1

Molecular Location: base pairs 21,797,389 to 21,941,869 on chromosome 12 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- ABC37
- ABCC9_HUMAN
- ATFB12
- ATP-binding cassette sub-family C member 9
- ATP-binding cassette sub-family C member 9 isoform SUR2A
- ATP-binding cassette sub-family C member 9 isoform SUR2B
- ATP-binding cassette transporter sub-family C member 9
- ATP-binding cassette, sub-family C (CFTR/MRP), member 9
- CANTU
- CMD10
- sulfonylurea receptor 2
- SUR2

Additional Information & Resources

Educational Resources

- Biochemistry (fifth edition, 2002): Specific Channels Can Rapidly Transport Ions Across Membranes
<https://www.ncbi.nlm.nih.gov/books/NBK22509/>
- The Human ATP-Binding Cassette (ABC) Transporter Superfamily: ABCC Genes
<https://www.ncbi.nlm.nih.gov/books/NBK3/#A191>

GeneReviews

- Dilated Cardiomyopathy Overview
<https://www.ncbi.nlm.nih.gov/books/NBK1309>

Genetic Testing Registry

- GTR: Genetic tests for ABCC9
<https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=10060%5Bgeneid%5D>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28ABCC9%5BTIAB%5D%29+OR+%28%28SUR2%5BTIAB%5D%29+OR+%28CMD1O%5BTIAB%5D%29+OR+%28sulfonylurea+receptor+2%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

OMIM

- ATP-BINDING CASSETTE, SUBFAMILY C, MEMBER 9
<http://omim.org/entry/601439>
- CARDIOMYOPATHY, DILATED, 10
<http://omim.org/entry/608569>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_ABCC9.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=ABCC9%5Bgene%5D>
- HGNC Gene Family: ATP binding cassette subfamily C
<http://www.genenames.org/cgi-bin/genefamilies/set/807>

- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=60
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/10060>
- UniProt
<http://www.uniprot.org/uniprot/O60706>

Sources for This Summary

- Bienengraeber M, Olson TM, Selivanov VA, Kathmann EC, O'Coilain F, Gao F, Karger AB, Ballew JD, Hodgson DM, Zingman LV, Pang YP, Alekseev AE, Terzic A. ABCC9 mutations identified in human dilated cardiomyopathy disrupt catalytic KATP channel gating. *Nat Genet.* 2004 Apr;36(4):382-7. Epub 2004 Mar 21.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15034580>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1995438/>
- Bryan J, Muñoz A, Zhang X, Düfer M, Drews G, Krippeit-Drews P, Aguilar-Bryan L. ABCC8 and ABCC9: ABC transporters that regulate K⁺ channels. *Pflugers Arch.* 2007 Feb;453(5):703-18. Epub 2006 Aug 8. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16897043>
- Harakalova M, van Harssel JJ, Terhal PA, van Lieshout S, Duran K, Renkens I, Amor DJ, Wilson LC, Kirk EP, Turner CL, Shears D, Garcia-Minaur S, Lees MM, Ross A, Venselaar H, Vriend G, Takanari H, Rook MB, van der Heyden MA, Asselbergs FW, Breur HM, Swinkels ME, Scurr IJ, Smithson SF, Knoers NV, van der Smagt JJ, Nijman IJ, Kloosterman WP, van Haelst MM, van Haaften G, Cuppen E. Dominant missense mutations in ABCC9 cause Cantú syndrome. *Nat Genet.* 2012 May 18;44(7):793-6. doi: 10.1038/ng.2324.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22610116>
- Solbach TF, König J, Fromm MF, Zolk O. ATP-binding cassette transporters in the heart. *Trends Cardiovasc Med.* 2006 Jan;16(1):7-15. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16387624>
- van Bon BW, Gilissen C, Grange DK, Hennekam RC, Kayserili H, Engels H, Reutter H, Ostergaard JR, Morava E, Tsiakas K, Isidor B, Le Merrer M, Eser M, Wieskamp N, de Vries P, Steehouwer M, Veltman JA, Robertson SP, Brunner HG, de Vries BB, Hoischen A. Cantú syndrome is caused by mutations in ABCC9. *Am J Hum Genet.* 2012 Jun 8;90(6):1094-101. doi: 10.1016/j.ajhg.2012.04.014. Epub 2012 May 17.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22608503>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3370286/>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/ABCC9>

Reviewed: January 2013
Published: February 14, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services