



## ADSL gene

adenylosuccinate lyase

### Normal Function

The *ADSL* gene provides instructions for making an enzyme called adenylosuccinate lyase. This enzyme performs two steps in the process that produces (synthesizes) purine nucleotides. These nucleotides are building blocks of DNA, its chemical cousin RNA, and molecules such as ATP that serve as energy sources in the cell. Adenylosuccinate lyase and other enzymes involved in purine synthesis form a group of proteins (a protein complex) called the purinosome. This complex comes together when there is a shortage of purines or when a large amount of purines is needed, such as during cell division. As part of this complex, adenylosuccinate lyase converts a molecule called succinylaminoimidazole carboxamide ribotide (SAICAR) to aminoimidazole carboxamide ribotide (AICAR) and converts succinyladenosine monophosphate (SAMP) to adenosine monophosphate (AMP).

### Health Conditions Related to Genetic Changes

#### adenylosuccinate lyase deficiency

More than 50 mutations in the *ADSL* gene have been found to cause adenylosuccinate lyase deficiency. This condition causes brain dysfunction (encephalopathy) that leads to delayed development of mental and motor skills (psychomotor delay), autistic behaviors that affect communication and social interaction, and seizures. Most of the mutations involved in this condition change single protein building blocks (amino acids) in the adenylosuccinate lyase enzyme. The altered enzymes have two to 20 percent of normal function, and studies suggest they are less able to form stable purinosomes.


A reduction of adenylosuccinate lyase function, possibly due to a shortage of purinosomes, leads to buildup of SAICAR and SAMP. These substances are converted through a different reaction to succinylaminoimidazole carboxamide riboside (SAICAr) and succinyladenosine (S-Ado). Detection of these substances in body fluids can help with diagnosis of adenylosuccinate lyase deficiency. Researchers believe that SAICAr and S-Ado are toxic; damage to brain tissue caused by one or both of these substances likely underlies the neurological problems that occur in adenylosuccinate lyase deficiency.

Studies suggest that the amount of SAICAr relative to S-Ado reflects the severity of adenylosuccinate lyase deficiency. Individuals with more SAICAr than S-Ado have more severe encephalopathy and psychomotor delay.

## Chromosomal Location

Cytogenetic Location: 22q13.1, which is the long (q) arm of chromosome 22 at position 13.1

Molecular Location: base pairs 40,346,500 to 40,387,408 on chromosome 22 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

## Other Names for This Gene

- adenylosuccinase
- adenylosuccinate lyase isoform a
- adenylosuccinate lyase isoform b
- AMPS
- ASASE
- ASL

## Additional Information & Resources

### Educational Resources

- Biochemistry (fifth edition, 2002): Purine Bases Can Be Synthesized de Novo or Recycled by Salvage Pathways  
<https://www.ncbi.nlm.nih.gov/books/NBK22385/>

### Genetic Testing Registry

- GTR: Genetic tests for ADSL  
<https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=158%5Bgeneid%5D>

## Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28ADSL%5BTIAB%5D%29+OR+%28adenylosuccinate+lyase%5BTIAB%5D%29%29+OR+%28adenylosuccinase%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

## OMIM

- ADENYLOSUCCINATE LYASE  
<http://omim.org/entry/608222>

## Research Resources

- ClinVar  
<https://www.ncbi.nlm.nih.gov/clinvar?term=ADSL%5Bgene%5D>
- HGNC Gene Symbol Report  
[http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?q=data/hgnc\\_data.php&hgnc\\_id=291](http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=291)
- NCBI Gene  
<https://www.ncbi.nlm.nih.gov/gene/158>
- UniProt  
<http://www.uniprot.org/uniprot/P30566>

## **Sources for This Summary**

- OMIM: ADENYLOSUCCINATE LYASE  
<http://omim.org/entry/608222>
- Ariyananda Lde Z, Lee P, Antonopoulos C, Colman RF. Biochemical and biophysical analysis of five disease-associated human adenylosuccinate lyase mutants. *Biochemistry*. 2009 Jun 16;48(23):5291-302. doi: 10.1021/bi802321m.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/19405474>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2745324/>
- Baresova V, Skopova V, Sikora J, Patterson D, Sovova J, Zikanova M, Kmoch S. Mutations of ATIC and ADSL affect purinosome assembly in cultured skin fibroblasts from patients with AICA-ribosiduria and ADSL deficiency. *Hum Mol Genet*. 2012 Apr 1;21(7):1534-43. doi: 10.1093/hmg/ddr591. Epub 2011 Dec 16.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/22180458>

- Ray SP, Deaton MK, Capodagli GC, Calkins LA, Sawle L, Ghosh K, Patterson D, Pegan SD. Structural and biochemical characterization of human adenylosuccinate lyase (ADSL) and the R303C ADSL deficiency-associated mutation. *Biochemistry*. 2012 Aug 21;51(33):6701-13. doi: 10.1021/bi300796y. Epub 2012 Aug 7.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/22812634>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3424377/>
  - Zikanova M, Skopova V, Hnizda A, Krijt J, Kmoch S. Biochemical and structural analysis of 14 mutant adsl enzyme complexes and correlation to phenotypic heterogeneity of adenylosuccinate lyase deficiency. *Hum Mutat*. 2010 Apr;31(4):445-55. doi: 10.1002/humu.21212.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/20127976>
- 

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/gene/ADSL>

Reviewed: December 2014

Published: February 14, 2017

Lister Hill National Center for Biomedical Communications  
U.S. National Library of Medicine  
National Institutes of Health  
Department of Health & Human Services