



CACNA1F gene

calcium voltage-gated channel subunit alpha1 F

Normal Function

The *CACNA1F* gene belongs to a family of genes that provide instructions for making calcium channels. These channels, which transport positively charged calcium atoms (calcium ions) across cell membranes, play a key role in a cell's ability to generate and transmit electrical signals.

The *CACNA1F* gene provides instructions for making one part (the alpha-1 subunit) of a calcium channel called CaV1.4. This subunit forms the hole (pore) in the cell membrane through which calcium ions can flow. CaV1.4 channels are found in many types of cells, although they play a particularly important role in a specialized tissue at the back of the eye called the retina. Within the retina, the channels are located in light-detecting cells called photoreceptors. The retina contains two types of photoreceptors: rods and cones. Rods are responsible for vision in low light. Cones provide vision in bright light, including color vision.

CaV1.4 channels appear to play a critical role in normal vision. Studies suggest they help relay visual signals from rods and cones to other retinal cells called bipolar cells. This signaling is an essential step in the transmission of visual information from the eyes to the brain.

Health Conditions Related to Genetic Changes

X-linked congenital stationary night blindness

More than 70 mutations in the *CACNA1F* gene have been identified in people with X-linked congenital stationary night blindness. Mutations in this gene are responsible for the incomplete form of the disorder, which is characterized by vision problems including a loss of sharpness (reduced acuity), nearsightedness (myopia), involuntary movements of the eyes (nystagmus), and eyes that do not look in the same direction (strabismus). Many affected individuals also have difficulty seeing in low light (night blindness).

CACNA1F mutations change the structure of the alpha-1 subunit, which alters or eliminates the function of CaV1.4 channels. These changes prevent the normal transport of calcium ions across the cell membrane of photoreceptor cells. A loss of calcium ion transport disrupts the ability of both rods and cones to transmit visual signals, which impairs vision.

Cone-rod dystrophy

Other disorders

Mutations in the *CACNA1F* gene cause several other rare disorders that impair vision. These include Åland Island eye disease, X-linked cone-rod dystrophy, an X-linked retinal disorder in New Zealand, and retinal and optic disc atrophy. Each of these disorders has been reported in only a few individuals or families worldwide. They cause vision problems similar to those of X-linked congenital stationary night blindness.

Researchers have identified at least one *CACNA1F* mutation that can cause Åland Island eye disease (also known as Forsius-Eriksson syndrome). This condition was first described in a family from the Åland Islands, which are in the Baltic Sea off the coast of Sweden. Åland Island eye disease is characterized by reduced visual acuity, nystagmus, an irregular curvature of the front part of the eye (astigmatism), myopia, abnormal color vision, and night blindness. The mutation associated with this disorder deletes a segment of genetic material from the *CACNA1F* gene, which significantly alters the structure of the alpha-1 subunit of CaV1.4 channels. These changes prevent the normal transport of calcium ions across the cell membrane of photoreceptor cells. A loss of calcium ion transport disrupts the ability of both rods and cones to transmit visual signals.

At least one other *CACNA1F* mutation is responsible for X-linked cone-rod dystrophy (also known as CORDX3). The signs and symptoms of this condition include reduced visual acuity, an increased sensitivity to light (photophobia), myopia, and impaired color vision. These vision problems tend to worsen over time. The mutation associated with this disorder deletes part of the alpha-1 subunit, which likely prevents the production of functional CaV1.4 channels. A loss of these channels keeps photoreceptor cells from relaying visual signals normally, which leads to impaired vision.

A *CACNA1F* mutation has also been found to cause an X-linked retinal disorder in a large Maori family from New Zealand. The major features of this disorder include reduced visual acuity, abnormal color vision, photophobia, and mild nystagmus. Some affected individuals have also had intellectual disability. The *CACNA1F* mutation associated with this condition alters a single protein building block (amino acid) in the alpha-1 subunit, which appears to overactivate CaV1.4 channels. The resulting increase in calcium ion transport probably disrupts the transmission of visual signals in the retina.


Another eye disorder, known as retinal and optic disc atrophy, has been associated with a *CACNA1F* mutation in two Japanese brothers. The affected individuals experienced a progressive decline in visual acuity and color vision. These vision problems were caused by deterioration of the retina, including an area called the optic disc (which is where the retina connects with the nerve that relays visual information to the brain). The *CACNA1F* mutation responsible for retinal and optic disc atrophy alters the structure of the alpha-1 subunit, which probably leads to the production of nonfunctional CaV1.4 channels. This mutation has also been identified

in at least one Japanese family with X-linked congenital stationary night blindness. It is unclear why this single genetic change can cause different vision abnormalities in different families.

Chromosomal Location

Cytogenetic Location: Xp11.23, which is the short (p) arm of the X chromosome at position 11.23

Molecular Location: base pairs 49,205,063 to 49,233,404 on the X chromosome (Homo sapiens Updated Annotation Release 109.20200522, GRCh38.p13) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- AIED
- CAC1F_HUMAN
- calcium channel, voltage-dependent, L type, alpha 1F subunit
- Cav1.4
- Cav1.4alpha1
- COD3
- COD4
- CORDX
- CORDX3
- CSNB2
- CSNB2A
- CSNBX2
- JM8
- JMC8
- OA2

Additional Information & Resources

Educational Resources

- Madame Curie Bioscience Database: α 1F/ CaV1.4
<https://www.ncbi.nlm.nih.gov/books/NBK6181/#A30881>
- Neuromuscular Disease Center, Washington University: Calcium channels
<https://neuromuscular.wustl.edu/mother/chan.html#ca>

Clinical Information from GeneReviews

- X-Linked Congenital Stationary Night Blindness
<https://www.ncbi.nlm.nih.gov/books/NBK1245>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28CACNA1F%5BTIAB%5D%29+OR+%28Cav1.4%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+2160+days%22%5Bdp%5D>

Catalog of Genes and Diseases from OMIM

- ALAND ISLAND EYE DISEASE
<http://omim.org/entry/300600>
- CALCIUM CHANNEL, VOLTAGE-DEPENDENT, ALPHA-1F SUBUNIT
<http://omim.org/entry/300110>
- CONE-ROD DYSTROPHY, X-LINKED, 3
<http://omim.org/entry/300476>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_CACNA1F.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=CACNA1F%5Bgene%5D>
- HGNC Gene Symbol Report
https://www.genenames.org/data/gene-symbol-report#!/hgnc_id/HGNC:1393
- Monarch Initiative
<https://monarchinitiative.org/gene/NCBIGene:778>
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/778>
- UniProt
<https://www.uniprot.org/uniprot/O60840>

Sources for This Summary

- Bech-Hansen NT, Naylor MJ, Maybaum TA, Pearce WG, Koop B, Fishman GA, Mets M, Musarella MA, Boycott KM. Loss-of-function mutations in a calcium-channel alpha1-subunit gene in Xp11.23 cause incomplete X-linked congenital stationary night blindness. *Nat Genet.* 1998 Jul;19(3):264-7. *Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/9662400>
- Boycott KM, Maybaum TA, Naylor MJ, Weleber RG, Robitaille J, Miyake Y, Bergen AA, Pierpont ME, Pearce WG, Bech-Hansen NT. A summary of 20 CACNA1F mutations identified in 36 families with incomplete X-linked congenital stationary night blindness, and characterization of splice variants. *Hum Genet.* 2001 Feb;108(2):91-7. *Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/11281458>
- Doering CJ, Peloquin JB, McRory JE. The Ca(v)1.4 calcium channel: more than meets the eye. *Channels (Austin).* 2007 Jan-Feb;1(1):3-10. Epub 2007 Jan 1. Review. *Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/19151588>
- Hemara-Wahanui A, Berjukow S, Hope CI, Dearden PK, Wu SB, Wilson-Wheeler J, Sharp DM, Lundon-Treweek P, Clover GM, Hoda JC, Striessnig J, Marksteiner R, Hering S, Maw MA. A CACNA1F mutation identified in an X-linked retinal disorder shifts the voltage dependence of Cav1.4 channel activation. *Proc Natl Acad Sci U S A.* 2005 May 24;102(21):7553-8. Epub 2005 May 16. *Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/15897456>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1140436/>
- Hoda JC, Zaghetto F, Koschak A, Striessnig J. Congenital stationary night blindness type 2 mutations S229P, G369D, L1068P, and W1440X alter channel gating or functional expression of Ca(v)1.4 L-type Ca²⁺ channels. *J Neurosci.* 2005 Jan 5;25(1):252-9. *Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/15634789>
- Hoda JC, Zaghetto F, Singh A, Koschak A, Striessnig J. Effects of congenital stationary night blindness type 2 mutations R508Q and L1364H on Cav1.4 L-type Ca²⁺ channel function and expression. *J Neurochem.* 2006 Mar;96(6):1648-58. Epub 2006 Feb 10. *Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/16476079>
- Hope CI, Sharp DM, Hemara-Wahanui A, Sissingh JI, Lundon P, Mitchell EA, Maw MA, Clover GM. Clinical manifestations of a unique X-linked retinal disorder in a large New Zealand family with a novel mutation in CACNA1F, the gene responsible for CSNB2. *Clin Experiment Ophthalmol.* 2005 Apr;33(2):129-36. *Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/15807819>
- Jalkanen R, Bech-Hansen NT, Tobias R, Sankila EM, Mäntyjärvi M, Forsius H, de la Chapelle A, Alitalo T. A novel CACNA1F gene mutation causes Aland Island eye disease. *Invest Ophthalmol Vis Sci.* 2007 Jun;48(6):2498-502. *Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/17525176>
- Jalkanen R, Mäntyjärvi M, Tobias R, Isosomppi J, Sankila EM, Alitalo T, Bech-Hansen NT. X linked cone-rod dystrophy, CORDX3, is caused by a mutation in the CACNA1F gene. *J Med Genet.* 2006 Aug;43(8):699-704. Epub 2006 Feb 27. *Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/16505158>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2564595/>
- McRory JE, Hamid J, Doering CJ, Garcia E, Parker R, Hamming K, Chen L, Hildebrand M, Beedle AM, Feldcamp L, Zamponi GW, Snutch TP. The CACNA1F gene encodes an L-type calcium channel with unique biophysical properties and tissue distribution. *J Neurosci.* 2004 Feb 18;24(7):1707-18. *Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/14973233>

- Nakamura M, Ito S, Piao CH, Terasaki H, Miyake Y. Retinal and optic disc atrophy associated with a CACNA1F mutation in a Japanese family. Arch Ophthalmol. 2003 Jul;121(7):1028-33.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12860808>
 - Peloquin JB, Rehak R, Doering CJ, McRory JE. Functional analysis of congenital stationary night blindness type-2 CACNA1F mutations F742C, G1007R, and R1049W. Neuroscience. 2007 Dec 5; 150(2):335-45. Epub 2007 Sep 14.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17949918>
 - Strom TM, Nyakatura G, Apfelstedt-Sylla E, Hellebrand H, Lorenz B, Weber BH, Wutz K, Gutwillinger N, Rütger K, Drescher B, Sauer C, Zrenner E, Meitinger T, Rosenthal A, Meindl A. An L-type calcium-channel gene mutated in incomplete X-linked congenital stationary night blindness. Nat Genet. 1998 Jul;19(3):260-3.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/9662399>
 - Wutz K, Sauer C, Zrenner E, Lorenz B, Alitalo T, Broghammer M, Hergersberg M, de la Chapelle A, Weber BH, Wissinger B, Meindl A, Pusch CM. Thirty distinct CACNA1F mutations in 33 families with incomplete type of XLCSNB and Cacna1f expression profiling in mouse retina. Eur J Hum Genet. 2002 Aug;10(8):449-56.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12111638>
-

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/CACNA1F>

Reviewed: May 2009

Published: August 17, 2020

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services