



FTL gene

ferritin light chain

Normal Function

The *FTL* gene provides instructions for making the ferritin light chain, which is one part (subunit) of a protein called ferritin. Ferritin is made up of 24 subunits formed into a hollow spherical molecule. The 24 subunits consist of varying numbers of the ferritin light chain and another subunit called the ferritin heavy chain, which is produced from another gene. The proportion of the two subunits varies in different tissues.

Ferritin stores and releases iron in cells. Each ferritin molecule can hold as many as 4,500 iron atoms inside its spherical structure. This storage capacity allows ferritin to regulate the amount of iron in cells and tissues. Iron is needed for the body to produce red blood cells.

Health Conditions Related to Genetic Changes

Hyperferritinemia-cataract syndrome

At least 31 mutations in the *FTL* gene have been identified in people with hyperferritinemia-cataract syndrome. Individuals affected by this disorder have an excess of ferritin in the blood (hyperferritinemia) and tissues of the body. A buildup of this protein begins early in life, leading to clouding of the lenses of the eyes (cataracts) in infancy.

The mutations that cause hyperferritinemia-cataract syndrome are found in a segment of the gene called the iron responsive element (IRE). The IRE normally can attach (bind) to a protein called the iron regulatory protein (IRP). When this binding occurs, the activity (expression) of the *FTL* gene is stopped to prevent too much ferritin light chain from being produced. This normally occurs when iron levels are low, because under those circumstances less ferritin is needed to store the iron. Mutations in the IRE segment of the *FTL* gene prevent it from binding with IRP, interfering with the mechanism by which ferritin production is matched to iron levels and resulting in excess ferritin being formed.

Neuroferritinopathy

At least four mutations in the *FTL* gene have been identified in people with neuroferritinopathy, a disorder in which iron gradually accumulates in the brain. These mutations all affect an area of the gene known as exon 4. The most common mutation detected in people with this disorder inserts the DNA building block (nucleotide) adenine between positions 460 and 461 in the gene sequence. Researchers believe that most families with this mutation descend from a common


ancestor who lived in northwest England before 1800. Three other known mutations are each found in a single individual or family affected by neuroferritinopathy.

Mutations in the *FTL* gene that cause neuroferritinopathy are believed to reduce ferritin's ability to store iron, resulting in the release of excess iron in nerve cells (neurons) of the brain. The cells may respond by producing more ferritin in an attempt to handle the free iron. Excess iron and ferritin accumulate in the brain, particularly in certain regions that help to control movement (basal ganglia), resulting in the movement problems and other neurological changes seen in neuroferritinopathy.

Chromosomal Location

Cytogenetic Location: 19q13.33, which is the long (q) arm of chromosome 19 at position 13.33

Molecular Location: base pairs 48,963,941 to 48,966,879 on chromosome 19 (Homo sapiens Updated Annotation Release 109.20190905, GRCh38.p13) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- ferritin L-chain
- ferritin L subunit
- ferritin light polypeptide-like 3
- ferritin, light polypeptide
- FRIL_HUMAN
- L apoferritin
- MGC71996
- NBIA3

Additional Information & Resources

Educational Resources

- Biochemistry (fifth edition, 2002): Genes Associated with Iron Metabolism are Translationally Regulated in Animals
<https://www.ncbi.nlm.nih.gov/books/NBK22400/#A4491>

Clinical Information from GeneReviews

- Neuroferritinopathy
<https://www.ncbi.nlm.nih.gov/books/NBK1141>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28FTL%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

Catalog of Genes and Diseases from OMIM

- FERRITIN LIGHT CHAIN
<http://omim.org/entry/134790>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_FTL.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=FTL%5Bgene%5D>
- HGNC Gene Symbol Report
https://www.genenames.org/data/gene-symbol-report/#!/hgnc_id/HGNC:3999
- Monarch Initiative
<https://monarchinitiative.org/gene/NCBIGene:2512>
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/2512>
- UniProt
<https://www.uniprot.org/uniprot/P02792>

Sources for This Summary

- Burn J, Chinnery PF. Neuroferritinopathy. *Semin Pediatr Neurol*. 2006 Sep;13(3):176-81.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17101456>
- Cazzola M. Role of ferritin and ferroportin genes in unexplained hyperferritinaemia. *Best Pract Res Clin Haematol*. 2005 Jun;18(2):251-63. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15737888>

- Craig JE, Clark JB, McLeod JL, Kirkland MA, Grant G, Elder JE, Toohey MG, Kowal L, Savoia HF, Chen C, Roberts S, Wirth MG, Mackey DA. Hereditary hyperferritinemia-cataract syndrome: prevalence, lens morphology, spectrum of mutations, and clinical presentations. *Arch Ophthalmol*. 2003 Dec;121(12):1753-61.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14662596>
- Crompton DE, Chinnery PF, Fey C, Curtis AR, Morris CM, Kierstan J, Burt A, Young F, Coulthard A, Curtis A, Ince PG, Bates D, Jackson MJ, Burn J. Neuroferritinopathy: a window on the role of iron in neurodegeneration. *Blood Cells Mol Dis*. 2002 Nov-Dec;29(3):522-31.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12547246>
- Curtis AR, Fey C, Morris CM, Bindoff LA, Ince PG, Chinnery PF, Coulthard A, Jackson MJ, Jackson AP, McHale DP, Hay D, Barker WA, Markham AF, Bates D, Curtis A, Burn J. Mutation in the gene encoding ferritin light polypeptide causes dominant adult-onset basal ganglia disease. *Nat Genet*. 2001 Aug;28(4):350-4.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11438811>
- OMIM: FERRITIN LIGHT CHAIN
<http://omim.org/entry/134790>
- Koorts AM, Viljoen M. Ferritin and ferritin isoforms I: Structure-function relationships, synthesis, degradation and secretion. *Arch Physiol Biochem*. 2007 Feb;113(1):30-54. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17522983>
- Levi S, Cozzi A, Arosio P. Neuroferritinopathy: a neurodegenerative disorder associated with L-ferritin mutation. *Best Pract Res Clin Haematol*. 2005 Jun;18(2):265-76. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15737889>
- Madsen E, Gitlin JD. Copper and iron disorders of the brain. *Annu Rev Neurosci*. 2007;30:317-37. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17367269>
- Millonig G, Muckenthaler MU, Mueller S. Hyperferritinaemia-cataract syndrome: worldwide mutations and phenotype of an increasingly diagnosed genetic disorder. *Hum Genomics*. 2010 Apr;4(4):250-62. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/20511138>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3525215/>
- Vidal R, Ghetti B, Takao M, Brefel-Courbon C, Uro-Coste E, Glazier BS, Siani V, Benson MD, Calvas P, Miravalle L, Rascol O, Delisle MB. Intracellular ferritin accumulation in neural and extraneural tissue characterizes a neurodegenerative disease associated with a mutation in the ferritin light polypeptide gene. *J Neuropathol Exp Neurol*. 2004 Apr;63(4):363-80.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15099026>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/FTL>

Reviewed: August 2012
Published: November 12, 2019

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services