



IFT80 gene

intraflagellar transport 80

Normal Function

The *IFT80* gene provides instructions for making a protein that is part of a group (complex) called IFT complex B. This complex is found in cell structures known as cilia. Cilia are microscopic, finger-like projections that stick out from the surface of cells. IFT complex B is involved in a process called intraflagellar transport (IFT), by which materials are carried within cilia. Specifically, this complex helps transport materials from the base of cilia to the tip.

IFT is essential for the assembly and maintenance of cilia. These cell structures play central roles in many different chemical signaling pathways, including a series of reactions called the Sonic Hedgehog pathway. These pathways are important for the growth and division (proliferation) and maturation (differentiation) of cells. In particular, Sonic Hedgehog appears to be essential for the proliferation and differentiation of cells that ultimately give rise to cartilage and bone.

Health Conditions Related to Genetic Changes

Asphyxiating thoracic dystrophy


Mutations in the *IFT80* gene were the first genetic changes found to cause asphyxiating thoracic dystrophy, an inherited disorder of bone growth characterized by a small chest, short ribs, and shortened bones in the arms and legs. At least six mutations in the *IFT80* gene have since been associated with this disorder. Most of these mutations change single protein building blocks (amino acids) in the IFT80 protein. IFT complex B made with the altered protein cannot function normally, which disrupts the transport of materials within cilia. Researchers speculate that these changes in IFT alter certain signaling pathways, including the Sonic Hedgehog pathway, which may underlie the abnormalities of bone growth characteristic of asphyxiating thoracic dystrophy.

In some affected individuals, asphyxiating thoracic dystrophy is also associated with abnormalities of the kidneys, liver, retinas, and other tissues. However, when the disorder results from *IFT80* gene mutations, its features are usually limited to problems with bone growth. The reasons for this difference are unknown.

Chromosomal Location

Cytogenetic Location: 3q25.33, which is the long (q) arm of chromosome 3 at position 25.33

Molecular Location: base pairs 160,256,986 to 160,399,225 on chromosome 3 (Homo sapiens Updated Annotation Release 109.20200522, GRCh38.p13) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- ATD2
- IFT80_HUMAN
- intraflagellar transport 80 homolog (Chlamydomonas)
- KIAA1374
- MGC126543
- WD repeat-containing protein 56
- WD repeat domain 56
- WDR56

Additional Information & Resources

Educational Resources

- Molecular Cell Biology (fourth edition, 2000): Cilia and Flagella: Structure and Movement
<https://www.ncbi.nlm.nih.gov/books/NBK21698/>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28IFT80%5BTI%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D>

Catalog of Genes and Diseases from OMIM

- INTRAFLAGELLAR TRANSPORT 80
<http://omim.org/entry/611177>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_IFT80.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=IFT80%5Bgene%5D>
- HGNC Gene Symbol Report
https://www.genenames.org/data/gene-symbol-report/#!/hgnc_id/HGNC:29262
- Monarch Initiative
<https://monarchinitiative.org/gene/NCBIGene:57560>
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/57560>
- UniProt
<https://www.uniprot.org/uniprot/Q9P2H3>

Sources for This Summary

- Beales PL, Bland E, Tobin JL, Bacchelli C, Tuysuz B, Hill J, Rix S, Pearson CG, Kai M, Hartley J, Johnson C, Irving M, Elcioglu N, Winey M, Tada M, Scambler PJ. IFT80, which encodes a conserved intraflagellar transport protein, is mutated in Jeune asphyxiating thoracic dystrophy. *Nat Genet.* 2007 Jun;39(6):727-9. Epub 2007 Apr 29.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17468754>
- Wang C, Yuan X, Yang S. IFT80 is essential for chondrocyte differentiation by regulating Hedgehog and Wnt signaling pathways. *Exp Cell Res.* 2013 Mar 10;319(5):623-32. doi: 10.1016/j.yexcr.2012.12.028. Epub 2013 Jan 16.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/23333501>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3908790/>
- Yang S, Wang C. The intraflagellar transport protein IFT80 is required for cilia formation and osteogenesis. *Bone.* 2012 Sep;51(3):407-17. doi: 10.1016/j.bone.2012.06.021. Epub 2012 Jul 4.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22771375>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3412883/>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/IFT80>

Reviewed: May 2015

Published: August 17, 2020

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services