



## IL17RC gene

interleukin 17 receptor C

### Normal Function

The *IL17RC* gene provides instructions for making a protein that is involved in immune system function, specifically in the body's defense against a fungus called *Candida*. When the immune system recognizes *Candida*, it generates cells called Th17 cells. These cells produce signaling molecules (cytokines) called the interleukin-17 (IL-17) family as part of an immune process called the IL-17 pathway. The IL-17 pathway creates inflammation, sending other cytokines and white blood cells that fight foreign invaders and promote tissue repair. In addition, the IL-17 pathway promotes the production of certain antimicrobial protein segments (peptides) that control growth of *Candida* on the surface of mucous membranes.

The protein produced from the *IL17RC* gene is present in many tissue types in the body, and is involved in cell signaling as part of the IL-17 pathway. Together with the protein produced from the *IL17RA* gene, it forms one of several receptors for IL-17 cytokines. Receptor proteins have specific sites into which certain other proteins, called ligands, fit like keys into locks. Certain IL-17 cytokines attach to receptors containing the IL17RC and IL17RA proteins, triggering signals that promote inflammation and the defense against *Candida* infection.

### Health Conditions Related to Genetic Changes

#### Familial candidiasis


At least three mutations in the *IL17RC* gene have been identified in people with familial candidiasis, an inherited tendency to develop infections caused by the *Candida* fungus (commonly known as yeast infections). Most people with familial candidiasis have chronic yeast infections of the skin, nails, and mucous membranes (collectively called chronic mucocutaneous candidiasis) beginning in early childhood.

*IL17RC* gene mutations result in an absence of IL17RC receptor protein. Loss of this protein impairs binding of IL-17 cytokines and IL-17 signaling. This impaired signaling decreases production of the antimicrobial peptides and inflammatory molecules that are critical for defense against *Candida*, and results in the chronic infection that occurs in familial candidiasis.

## Chromosomal Location

Cytogenetic Location: 3p25.3, which is the short (p) arm of chromosome 3 at position 25.3

Molecular Location: base pairs 9,917,074 to 9,933,627 on chromosome 3 (Homo sapiens Annotation Release 109, GRCh38.p12) (NCBI)



Credit: Genome Decoration Page/NCBI

## Other Names for This Gene

- IL-17 receptor C
- IL-17RL
- IL17F receptor
- interleukin-17 receptor homolog
- interleukin-17 receptor-like protein

## Additional Information & Resources

### Educational Resources

- Immunobiology: The Immune System in Health and Disease (fifth edition, 2001): The Front Line of Host Defense  
<https://www.ncbi.nlm.nih.gov/books/NBK27105/>

## Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28IL17RC%5BTIAB%5D%29+OR+%28interleukin+17+receptor+C%5BTIAB%5D%29%29+OR+%28%28IL-17+receptor+C%5BTIAB%5D%29+OR+%28IL-17RL%5BTIAB%5D%29+OR+%28IL17F+receptor%5BTIAB%5D%29+OR+%28interleukin-17+receptor+C+isoform+1+precursor%5BTIAB%5D%29+OR+%28interleukin-17+receptor+C+isoform+2+precursor%5BTIAB%5D%29+OR+%28interleukin-17+receptor+C+isoform+3+precursor%5BTIAB%5D%29+OR+%28interleukin-17+receptor+C+isoform+4+precursor%5BTIAB%5D%29+OR+%28interleukin-17+receptor+C+isoform+5+precursor%5BTIAB%5D%29+OR+%28interleukin-17+receptor+C+isoform+6+precursor%5BTIAB%5D%29+OR+%28interleukin-17+receptor+homolog%5BTIAB%5D%29+OR+%28interleukin-17+receptor-like+protein%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

## Catalog of Genes and Diseases from OMIM

- INTERLEUKIN 17 RECEPTOR C  
<http://omim.org/entry/610925>

## Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology  
[http://atlasgeneticsoncology.org/Genes/GC\\_IL17RC.html](http://atlasgeneticsoncology.org/Genes/GC_IL17RC.html)
- ClinVar  
<https://www.ncbi.nlm.nih.gov/clinvar?term=IL17RC%5Bgene%5D>
- HGNC Gene Symbol Report  
[https://www.genenames.org/data/gene-symbol-report#!/hgnc\\_id/HGNC:18358](https://www.genenames.org/data/gene-symbol-report#!/hgnc_id/HGNC:18358)
- Monarch Initiative  
<https://monarchinitiative.org/gene/NCBIGene:84818>
- NCBI Gene  
<https://www.ncbi.nlm.nih.gov/gene/84818>
- UniProt  
<https://www.uniprot.org/uniprot/Q8NAC3>

## Sources for This Summary

- Ho AW, Shen F, Conti HR, Patel N, Childs EE, Peterson AC, Hernández-Santos N, Kolls JK, Kane LP, Ouyang W, Gaffen SL. IL-17RC is required for immune signaling via an extended SEF/IL-17R signaling domain in the cytoplasmic tail. *J Immunol*. 2010 Jul 15;185(2):1063-70. doi: 10.4049/jimmunol.0903739. Epub 2010 Jun 16.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/20554964>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2897912/>
- Huppler AR, Bishu S, Gaffen SL. Mucocutaneous candidiasis: the IL-17 pathway and implications for targeted immunotherapy. *Arthritis Res Ther*. 2012 Jul 23;14(4):217. doi: 10.1186/ar3893. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/22838497>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3580547/>
- OMIM: INTERLEUKIN 17 RECEPTOR C  
<http://omim.org/entry/610925>
- Ling Y, Cypowyj S, Aytekin C, Galicchio M, Camcioglu Y, Nepesov S, Ikinciogullari A, Dogu F, Belkadi A, Levy R, Migaud M, Boisson B, Bolze A, Itan Y, Goudin N, Cottineau J, Picard C, Abel L, Bustamante J, Casanova JL, Puel A. Inherited IL-17RC deficiency in patients with chronic mucocutaneous candidiasis. *J Exp Med*. 2015 May 4;212(5):619-31. doi: 10.1084/jem.20141065. Epub 2015 Apr 27.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/25918342>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4419340/>
- Plantinga TS, Johnson MD, Scott WK, Joosten LA, van der Meer JW, Perfect JR, Kullberg BJ, Netea MG. Human genetic susceptibility to *Candida* infections. *Med Mycol*. 2012 Nov;50(8):785-94. doi: 10.3109/13693786.2012.690902. Epub 2012 Jun 4. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/22662758>
- Puel A, Cypowyj S, Maródi L, Abel L, Picard C, Casanova JL. Inborn errors of human IL-17 immunity underlie chronic mucocutaneous candidiasis. *Curr Opin Allergy Clin Immunol*. 2012 Dec; 12(6):616-22. doi: 10.1097/ACI.0b013e328358cc0b. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/23026768>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3538358/>
- Smeekens SP, van de Veerdonk FL, Kullberg BJ, Netea MG. Genetic susceptibility to *Candida* infections. *EMBO Mol Med*. 2013 Jun;5(6):805-13. doi: 10.1002/emmm.201201678. Epub 2013 Apr 30. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/23629947>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3779444/>

---

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/gene/IL17RC>

Reviewed: September 2016

Published: June 11, 2019

Lister Hill National Center for Biomedical Communications  
U.S. National Library of Medicine  
National Institutes of Health  
Department of Health & Human Services