



TSC2 gene

TSC complex subunit 2

Normal Function

The *TSC2* gene provides instructions for producing a protein called tuberin, whose function is not fully understood. Within cells, tuberin interacts with a protein called hamartin, which is produced from the *TSC1* gene. These two proteins help control cell growth and size. Proteins that normally prevent cells from growing and dividing too fast or in an uncontrolled way are known as tumor suppressors. Hamartin and tuberin carry out their tumor suppressor function by interacting with and regulating a wide variety of other proteins.

Health Conditions Related to Genetic Changes

Lymphangiomyomatosis

Mutations in the *TSC2* gene cause most cases of a disorder called lymphangiomyomatosis (LAM). This destructive lung disease is characterized by the abnormal overgrowth of smooth muscle-like tissue in the lungs. It occurs almost exclusively in women, causing coughing, shortness of breath, chest pain, and lung collapse.

LAM can occur alone (isolated or sporadic LAM) or in combination with a condition called tuberous sclerosis complex (described below). Researchers suggest that sporadic LAM is caused by a random mutation in the *TSC2* gene that occurs very early in development. As a result, some of the body's cells have a normal version of the gene, while others have the mutated version. This situation is called mosaicism. When a mutation occurs in the other copy of the *TSC2* gene in certain cells during a woman's lifetime (a somatic mutation), she may develop LAM.

Tuberous sclerosis complex

More than 1,100 mutations in the *TSC2* gene have been identified in individuals with tuberous sclerosis complex, a condition characterized by developmental problems and the growth of noncancerous tumors in many parts of the body. Most of these mutations insert or delete a small number of DNA building blocks (base pairs) in the *TSC2* gene. Other mutations change a single base pair in the *TSC2* gene or create a premature stop signal in the instructions for making tuberin.

People with *TSC2*-related tuberous sclerosis complex are born with one mutated copy of the *TSC2* gene in each cell. This mutation prevents the cell from making functional tuberin from that copy of the gene. However, enough tuberin is usually produced from the other, normal copy of the *TSC2* gene to regulate cell growth


effectively. For some types of tumors to develop, a second mutation involving the other copy of the gene must occur in certain cells during a person's lifetime.

When both copies of the *TSC2* gene are mutated in a particular cell, that cell cannot produce any functional tuberlin. The loss of this protein allows the cell to grow and divide in an uncontrolled way to form a tumor. A shortage of tuberlin also interferes with the normal development of certain cells. In people with *TSC2*-related tuberous sclerosis complex, a second *TSC2* gene mutation typically occurs in multiple cells over an affected person's lifetime. The loss of tuberlin in different types of cells disrupts normal development and leads to the growth of tumors in many different organs and tissues.

Chromosomal Location

Cytogenetic Location: 16p13.3, which is the short (p) arm of chromosome 16 at position 13.3

Molecular Location: base pairs 2,047,804 to 2,089,491 on chromosome 16 (Homo sapiens Updated Annotation Release 109.20190905, GRCh38.p13) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- PPP1R160
- TSC2_HUMAN
- tuberlin
- tuberous sclerosis 2

Additional Information & Resources

Educational Resources

- Cancer Medicine (sixth edition, 2003): Tuberous Sclerosis
<https://www.ncbi.nlm.nih.gov/books/NBK13658/#A19617>

Clinical Information from GeneReviews

- Tuberous Sclerosis Complex
<https://www.ncbi.nlm.nih.gov/books/NBK1220>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28TSC2%5BTIAB%5D%29+OR+%28tuberous+sclerosis+2%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5BIa%5D+AND+human%5Bmh%5D+AND+%22last+720+days%22%5Bdp%5D>

Catalog of Genes and Diseases from OMIM

- TSC2 GENE
<http://omim.org/entry/191092>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
<http://atlasgeneticsoncology.org/Genes/TSC2ID184.html>
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=TSC2%5Bgene%5D>
- HGNC Gene Symbol Report
https://www.genenames.org/data/gene-symbol-report#!/hgnc_id/HGNC:12363
- Leiden Open Variation Database: TSC2 Homepage
http://chromium.lovd.nl/LOVD2/TSC/home.php?select_db=TSC2
- Monarch Initiative
<https://monarchinitiative.org/gene/NCBIGene:7249>
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/7249>
- UniProt
<https://www.uniprot.org/uniprot/P49815>

Sources for This Summary

- Carsillo T, Astrinidis A, Henske EP. Mutations in the tuberous sclerosis complex gene TSC2 are a cause of sporadic pulmonary lymphangiomyomatosis. *Proc Natl Acad Sci U S A*. 2000 May 23; 97(11):6085-90.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10823953>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC18562/>
- Chorianopoulos D, Stratakos G. Lymphangiomyomatosis and tuberous sclerosis complex. *Lung*. 2008 Jul-Aug;186(4):197-207. doi: 10.1007/s00408-008-9087-5. Epub 2008 Apr 12. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18408969>
- Goncharova EA, Krymskaya VP. Pulmonary lymphangiomyomatosis (LAM): progress and current challenges. *J Cell Biochem*. 2008 Feb 1;103(2):369-82. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17541983>

- Huang J, Manning BD. The TSC1-TSC2 complex: a molecular switchboard controlling cell growth. *Biochem J*. 2008 Jun 1;412(2):179-90. doi: 10.1042/BJ20080281. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18466115>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2735030/>
- Juvet SC, McCormack FX, Kwiatkowski DJ, Downey GP. Molecular pathogenesis of lymphangioliomyomatosis: lessons learned from orphans. *Am J Respir Cell Mol Biol*. 2007 Apr; 36(4):398-408. Epub 2006 Nov 10. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17099139>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2176113/>
- Mak BC, Yeung RS. The tuberous sclerosis complex genes in tumor development. *Cancer Invest*. 2004;22(4):588-603. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15565817>
- Martignoni G, Pea M, Reghellin D, Gobbo S, Zamboni G, Chilosi M, Bonetti F. Molecular pathology of lymphangioliomyomatosis and other perivascular epithelioid cell tumors. *Arch Pathol Lab Med*. 2010 Jan;134(1):33-40. doi: 10.1043/2008-0542-RAR1.1.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/20073603>
- Narayanan V. Tuberous sclerosis complex: genetics to pathogenesis. *Pediatr Neurol*. 2003 Nov; 29(5):404-9. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14684235>
- Rosner M, Hanneder M, Siegel N, Valli A, Hengstschläger M. The tuberous sclerosis gene products hamartin and tuberin are multifunctional proteins with a wide spectrum of interacting partners. *Mutat Res*. 2008 Mar-Apr;658(3):234-46. doi: 10.1016/j.mrrev.2008.01.001. Epub 2008 Jan 12. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18291711>
- Sampson JR. TSC1 and TSC2: genes that are mutated in the human genetic disorder tuberous sclerosis. *Biochem Soc Trans*. 2003 Jun;31(Pt 3):592-6.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12773162>
- Yeung RS. Multiple roles of the tuberous sclerosis complex genes. *Genes Chromosomes Cancer*. 2003 Dec;38(4):368-75. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14566857>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/TSC2>

Reviewed: June 2010

Published: October 1, 2019

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services